

10011337 PCT/PTO 15 FEB 2002

FORM PTO-1390 (REV 12-2001)		U.S. DEPARTMENT OF COMMERCE PATENT AND TRADEMARK OFFICE	ATTORNEY'S DOCKET NUMBER 514413-3915
TRANSMITTAL LETTER TO THE UNITED STATES DESIGNATED/ELECTED OFFICE (DO/EO/US) CONCERNING A FILING UNDER 35 U.S.C. 371			U.S. APPLICATION NO. (If known see 37 C.F.R. 1.5) 10/049731
INTERNATIONAL APPLICATION NO. PCT/EP00/09339	INTERNATIONAL FILING DATE 8 SEPTEMBER 2000	PRIORITY DATE CLAIMED 8 SEPTEMBER 1999	
TITLE OF INVENTION NEW HERBICIDAL COMPOSITIONS			
APPLICANT(S) FOR DO/EO/US Ken PALLETT and Ashley SLATER			
Applicants herewith submit to the United States Designated/Elected Office (DO/EO/US) the following items and other information:			
1. <input checked="" type="checkbox"/> This is a FIRST submission of items concerning a filing under 35 U.S.C. 371. 2. <input type="checkbox"/> This is a SECOND or SUBSEQUENT submission of items concerning a filing under 35 U.S.C. 371. 3. <input checked="" type="checkbox"/> This is an express request to promptly begin national examination procedures (35 U.S.C. 371(f)). The submission must include items (5), (6), (9) and (21) indicated below. 4. <input checked="" type="checkbox"/> The US has been elected by the expiration of 19 months from the priority date (PCT Article 31). 5. <input checked="" type="checkbox"/> A copy of the International Application as filed (35 U.S.C. 371(c)(2)). a. <input type="checkbox"/> is attached hereto (required only if not communicated by the International Bureau). b. <input checked="" type="checkbox"/> has been communicated by the International Bureau. c. <input type="checkbox"/> is not required, as the application was filed in the United States Receiving Office (RO/US). 6. <input checked="" type="checkbox"/> An English language translation of the International Application as filed (35 U.S.C. 371(c)(2)). a. <input checked="" type="checkbox"/> is attached hereto. b. <input type="checkbox"/> has been previously submitted under 35 U.S.C. 154(d)(4). 7. <input checked="" type="checkbox"/> Amendments to the claims of the International Application under PCT Article 19 (35 U.S.C. 371(c)(3)). a. <input type="checkbox"/> are attached hereto (required only if not communicated by the International Bureau). b. <input type="checkbox"/> have been communicated by the International Bureau. c. <input type="checkbox"/> have not been made; however, the time limit for making such amendments has NOT expired. d. <input checked="" type="checkbox"/> have not been made and will not be made. 8. <input type="checkbox"/> A English language translation of the amendments to the claims under PCT Article 19 (35 U.S.C. 371(c)(3)). 9. <input checked="" type="checkbox"/> An oath or declaration of the inventor(s) (35 U.S.C. 371(c)(4)). 10. <input type="checkbox"/> An English language translation of the annexes to the International Preliminary Examination Report under PCT Article 36 (35 U.S.C. 371(c)(5)). Items 11 to 20 below concern document(s) or information included: 11. <input checked="" type="checkbox"/> An Information Disclosure Statement under 37 CFR 1.97 and 1.98. 12. <input checked="" type="checkbox"/> An assignment document for recording. A separate cover sheet in compliance with 37 CFR 3.28 and 3.31 is included. 13. <input type="checkbox"/> A FIRST preliminary amendment 14. <input type="checkbox"/> A SECOND or SUBSEQUENT preliminary amendment. 15. <input type="checkbox"/> A substitute specification 16. <input type="checkbox"/> A change of power of attorney and/or address letter 17. <input type="checkbox"/> A computer-readable form of the sequence listing in accordance with PCT Rule 13ter.2 and 35 U.S.C. 1.821 - 1.825. 18. <input type="checkbox"/> A second copy of the published international application under 35 U.S.C. 154(d)(4) 19. <input type="checkbox"/> A second copy of the English language translation of the international application under 35 U.S.C. 154(d)(4). 20. <input checked="" type="checkbox"/> Other items or information: PCT/RO/101, PCT/IPEA/416, 409, PCT/IB/308, PCT/ISA/210			
		EXPRESS MAIL Mailing Label Number EV001580947US Date of Deposit: February 15, 2002 I hereby certify that this paper or fee is being deposited with the United States Postal Service "Express Mail Post Office to Addressee" Service under 37 CFR 1.10 on the date indicated above and is addressed to the Assistant Commissioner for Patents and Trademarks, Box PCT Washington, DC 20231 Charles Jackson (Typed or printed name of person mailing paper or fee) Charles Jackson (Signature of person mailing paper or fee)	

U.S. APPLICATION NO. (If known, see 37 C.F.R. 1.60) 10/049731		INTERNATIONAL APPLICATION NO. PCT/EP00/09339		ATTORNEY'S DOCKET NO. 514413-3915	
---	--	---	--	--------------------------------------	--

21. <input checked="" type="checkbox"/> The following fees are submitted				CALCULATIONS PTO USE ONLY	
BASIC NATIONAL FEE (37 CFR 1.492(a)(1)-(5): Neither international preliminary examination fee (37 CFR 1.482) nor international search fee (37 CFR 1.445(a)(2)) paid to USPTO and International Search Report not prepared by the EPO or JPO \$1040.00 International preliminary examination fee (37 C.F.R. 1.482) not paid to USPTO but International Search Report prepared by the EPO or JPO \$890.00 International preliminary examination fee (37 CFR 1.482) not paid to USPTO but international search fee (37 CFR 1.445(a)(2)) paid to USPTO \$740.00 International preliminary examination fee paid to USPTO (37 CFR 1.482) but all claims did not satisfy provisions of PCT Article 33(1)-(4)..... \$710.00 International preliminary examination fee paid to USPTO (37 CFR 1.482) and all claims satisfied provisions of PCT Article 33(1)-(4)..... \$100.00					
ENTER APPROPRIATE BASIC FEE AMOUNT =				\$ 890.00	
Surcharge of \$130.00 for furnishing the oath or declaration later than <input type="checkbox"/> 20 <input type="checkbox"/> 30 months from the earliest claimed priority date (37 CFR 1.492(e)).				\$	
CLAIMS	NUMBER FILED	NUMBER EXTRA	RATE	\$	
Total Claims	<u>22</u> - 20 =	<u>2</u>	x \$18.00	\$ 36.00	
Independent Claims	<u>3</u> - 3 =	<u>0</u>	x \$84.00	\$ 0.00	
MULTIPLE DEPENDENT CLAIM(S) (if applicable)				+ \$280.00	
TOTAL OF ABOVE CALCULATIONS =				\$	
<input type="checkbox"/> Applicant claims small entity status. See 37 C.F.R. 1.27. The fees indicated above are reduced by 1/2.				+	
SUBTOTAL =				\$ 926.00	
Processing fee of \$130.00 for furnishing the English translation later than <input type="checkbox"/> 20 <input type="checkbox"/> 30 months from the earliest claimed priority date (37 CFR 1.492(f))				\$	
TOTAL NATIONAL FEE =				\$	
Fee for recording the enclosed assignments (37 CFR 1.21(h)) The assignment must be accompanied by an appropriate cover sheet (37 CFR 3.28, 3.31) \$40.00 per property				+ \$ 40.00	
TOTAL FEES ENCLOSED =				\$ 966.00	
				Amount to be refunded:	
				\$	
				charged:	
				\$	

a. ☒ A check in the amount of **\$966.00** to cover the above fees is enclosed.

b. ☐ Please charge my Deposit Account No. _____ in the amount of \$ _____ to cover the above fees. A duplicate copy of this sheet is enclosed.

c. ☒ The Commissioner is hereby authorized to charge any additional fees which may be required, or credit any overpayment to Deposit Account No. **50-0320**. A duplicate copy of this sheet is enclosed.

d. ☐ Fees are to be charged to a credit card. WARNING: Information on this form may become public. **Credit card information should not be included on this form.** Provide credit card information and authorization on PTO-2038.

NOTE: Where an appropriate time limit under 37 CFR 1.494 or 1.495 has not been met, a petition to revive (37 CFR 1.137(a) or (b)) must be filed and granted to restore the application to pending status.

SEND ALL CORRESPONDENCE TO:

William S. Frommer, Esq.
Frommer Lawrence & Haug LLP
745 Fifth Avenue
New York, New York 10151

 SIGNATURE

 William F. Lawrence
 NAME

 28,029
 REGISTRATION NUMBER

Dated: February 15, 2002

New Herbicidal Compositions

Background of the Invention

The present invention relates to the safening of herbicidal compounds, in particular the safening of benzoylisoxazole and/or dione derivatives which are useful for the growing of crops in particular for maize (*Zea mays*) and to compositions useful for such treatment.

Discussion of Related Art

An important factor influencing the usefulness of a given herbicide is its selectivity toward crops. In some cases, a beneficial crop is susceptible to the effects of a herbicide when applied at application rates needed to control weed growth. In addition, certain herbicidal compounds are phytotoxic to some weed species but not to others. This may render such herbicides unsuitable for controlling weeds in the presence of certain crops. To be effective, a herbicide must cause minimal damage (preferably no damage) to the beneficial crop while maximising the damage to weed species which infest the locus of the crop. Reduction in herbicidal injury to crops without an unacceptable reduction in the herbicidal action can be accomplished by the use of crop protectants known as "antidotes" or "safeners".

Identification of an antidote which safens a herbicide in crops is a complicated task. The precise mechanism by which an antidote reduces herbicidal crop injury has not been established. In general, the safening effect of a compound is specific to the herbicidal partner and the crop where the active ingredients are applied. Benzoylisoxazoles are known to possess herbicidal properties for example, European Patent Publication Nos. 0418175, 0487357, 0527036 and 0560482. European Patent Publication Nos. 0496630, 0496631, 0625505 and 0625508 disclose certain dione derivatives possessing herbicidal properties. In general such herbicides are very active against broad-leaved and grass weeds by pre- and/or post-emergence application. The method of controlling vegetation with these compounds comprises applying a herbicidally effective amount of the compounds, usually with an inert carrier or diluent, to the area where herbicidal control is desired. However, the herbicidal benzoylisoxazole and/or dione compounds have been found in some instances to adversely affect or interfere with the development of crop plants, especially maize crops. The effective use of these herbicides for controlling weeds in the presence of such crops may be enhanced by, or may

require in certain instances, the addition of a compound which is antidotally effective with the herbicide.


Although it is possible to say in general terms that herbicides may be used in the presence of an antidote, the problem in identifying specific antidotes for specific crops at appropriate rates to control weed growth, is substantial.

The applicants have found that certain compounds, in particular when applied as described hereinafter, are effective antidotes for the protection of crops, especially maize crops, from herbicidal injury or the reduction of herbicidal injury caused by the application of an amount of a benzoylisoxazole and/or dione compound (optionally in admixture with a partner herbicide) effective to control the growth of weeds.

It is an object of the present invention to provide compositions of benzoylisoxazoles and/or dione herbicides in combination with antidotes, said compositions providing a reduction in crop injury, especially to maize (*Zea mays*), arising from the phytotoxicity of the herbicides, and a method of using said herbicides and antidotes.




Description of the Invention


The present invention provides a method of reducing phytotoxicity to a crop (especially maize) at a locus caused by the application thereto of a herbicidal benzoylisoxazole and/or dione derivative of formula (I):




wherein:

A is a group (A-1) to (A-7):







(A-4)



(A-5)




(A-6)




(A-7)

5 or a corresponding formula (A-6a) or (A-7a):



(A-6a)



(A-7a)

- in which the position of the carbonyl group and the group Q are reversed and the double bond in the ring is attached to the carbon atom attached to the group Q;
- 10 R is a hydrogen atom or a halogen atom; a straight- or branched chain alkyl, alkenyl or alkynyl group containing from one to six carbon atoms which is optionally substituted by one or more halogen atoms; a cycloalkyl group containing from 3 to 6 carbon atoms optionally substituted by one or more groups R^5 , one or more halogen atoms or a group $-CO_2R^3$; or a group selected from $-CO_2R^3$, $-COR^5$,
 15 cyano, nitro, $-CONR^3R^4$ and $-S(O)_kR^{13}$;

R^1 is a straight- or branched-chain alkyl, alkenyl or alkynyl group containing up to six carbon atoms which is optionally substituted by one or more halogen atoms; or a cycloalkyl group containing from three to six carbon atoms optionally substituted by one or more groups R^5 or one or more halogen atoms;

R^2 is a halogen atom; a straight- or branched-chain alkyl, alkenyl or alkynyl group containing up to six carbon atoms which is optionally substituted by one or more halogen atoms; a straight- or branched-chain alkyl group containing up to six carbon atoms which is substituted by one or more groups $-OR^5$; or a group
 5 selected from nitro, cyano, $-CO_2R^5$, $-S(O)_pR^6$, $-O(CH_2)_mOR^5$, $-COR^5$, $-NR^{11}R^{12}$, $-N(R^8)SO_2R^7$, $-N(R^8)CO_2R^7$, $-OR^5$, $-OSO_2R^7$, $-SO_2NR^3R^4$, $-CONR^3R^4$, $-CSNR^3R^4$, $-(CR^9R^{10})_y-S(O)_qR^7$ and $-SF_5$;

or two groups R^2 , on adjacent carbon atoms of the phenyl ring may, together with the carbon atoms to which they are attached, form a 5 to 7 membered saturated
 10 or unsaturated heterocyclic ring containing up to three ring heteroatoms selected from nitrogen, oxygen and sulfur, which ring is optionally substituted by one or more groups selected from halogen, nitro, $-S(O)_pR^{13}$, C_{1-4} alkyl, C_{1-4} alkoxy, C_{1-4} haloalkyl, C_{1-4} haloalkoxy, $=O$ (or a 5- or 6- membered cyclic acetal thereof), and $=NO-R^3$, it being understood that a sulphur atom, where present in the ring,
 15 may be in the form of a group $-SO-$ or $-SO_2-$;

z is an integer from one to five: when z is greater than one the groups R^2 may be the same or different;

R^3 , R^4 and R^{109} are each independently a hydrogen atom, or a straight- or branched chain alkyl group containing up to six carbon atoms which is optionally
 20 substituted by one or more halogen atoms;

R^5 and R^{110} are each independently a straight- or branched-chain alkyl group containing up to six carbon atoms which is optionally substituted by one or more halogen atoms or a straight- or branched-chain alkenyl or alkynyl group containing from two to six (preferably from three to six) carbon atoms which is optionally
 25 substituted by one or more halogen atoms;

R^6 and R^7 , which may be the same or different, are each R^5 ; or phenyl optionally substituted by from one to five groups which may be the same or different selected from a halogen atom, a straight- or branched-chain alkyl group containing up to six carbon atoms which is optionally substituted by one or more halogen
 30 atoms, nitro, cyano, $-CO_2R^5$, $-S(O)_pR^{13}$, $-NR^{11}NR^{12}$, $-OR^5$ and $-CONR^3R^4$;
 R^8 , R^9 and R^{10} are each a hydrogen atom or R^6 ;

R^{11} and R^{12} are each a hydrogen atom or R^5 ;

R^{13} and R^{111} are each a straight- or branched-chain alkyl group containing up to six carbon atoms which is optionally substituted by one or more halogen atoms;

Q is hydroxy, C1-6 alkoxy, OR^{112} , SR^{112} or SR^{111} ;

5 L is oxygen or NR^{109} ;

R^{14} , R^{14a} , R^{14b} , R^{15} , R^{15a} , R^{15b} , R^{16} , R^{16a} , R^{16b} , R^{100} , R^{101} , R^{102} , R^{103} , R^{104} , R^{105} , R^{106} , R^{107} and R^{108} are each the same or different groups selected from hydrogen, R^{110} , $-(CH_2)_uCO_2R^{109}$, halogen, cyano, C1-6 alkoxy,

10 $-(CH_2)_x$ -[phenyl optionally substituted by from one to five groups R^{113} which may be the same or different], and cycloalkyl containing from three to six carbon atoms optionally substituted by C1-6 alkyl or $-S(O)_pR^{111}$;

R^{112} is phenyl optionally substituted by from one to five groups selected from halogen, C1-6 alkyl, C1-6 haloalkyl, C1-6 alkoxy and nitro;

R^{113} is a group selected from halogen, R^{114} , nitro, cyano, $-CO_2R^{115}$,

15 $-S(O)_pR^{111}$, $-OR^{111}$ and $-NR^{115}R^{116}$;

R^{114} is a straight- or branched- chain alkyl group containing one to three carbon atoms optionally substituted by one or more halogen atoms;

R^{115} and R^{116} which may be the same or different, are each a hydrogen atom or R^{110} ;

20 p, q and u are each independently zero, one or two;

k and m are each one, two or three;

x is zero or one;

y is an integer from one to four; when y is greater than one, the groups R^9 and R^{10} may be the same or different;

25 or an agriculturally acceptable salt or metal complex thereof; which method comprises applying to the locus of the crop, preferably before the herbicidal compound, an antidotally effective amount of an antidote compound, and optional partner herbicide.

It will be understood that the said antidote is, in general, antidotally effective for 30 said benzoylisoxazole and/or dione derivative.

It will be understood that antidotes used in the method of the invention may form for example salts, and that the use of such salts is also embraced by the invention.

- In this patent specification including the accompanying claims it is understood that the term 'agriculturally acceptable salts' is meant salts the cations or anions of which are known and accepted in the art for the formation of salts for agricultural or horticultural use. Preferably the salts are water-soluble. Suitable salts with
- 5 bases include alkali metal (e.g. sodium and potassium), alkaline earth metal (e.g. calcium and magnesium), ammonium and amine (e.g. diethanolamine, triethanolamine, octylamine, morpholine and dioctylmethylamine) salts. Suitable acid addition salts, e.g. formed by compounds of formula (I) containing an amino group, include salts with inorganic acids, for example hydrochlorides, sulphates,
- 10 phosphates and nitrates and salts with organic acids for example acetic acid. It will be understood that the term 'dione' as used in this specification including the accompanying claims does not exclude the possible presence of additional C=O groups as in triones.
- By the term "metal complexes" is meant compounds in which one or both of
- 15 the oxygen atoms of the 1,3-dione of formula (A-3), (A-4), (A-5), (A-6), (A-7), (A-6a) or (A-7a) act as chelating agents to a metal cation. Examples of such cations include zinc, manganese, cupric, cuprous, ferric, ferrous, titanium and aluminium.
- 20 Compounds of formula (II) may exist in enolic tautomeric forms that may give rise to geometric isomers around the enolic double bond. Furthermore in certain cases the above substituents may contribute to optical isomerism and/or stereoisomerism. All such forms and mixtures thereof are embraced by the present invention.
- 25 It is to be understood that in this specification compounds comprising a cyclohexane ring corresponding to formula (A-6) or (A-7) or a precursor thereof include the compounds with the corresponding formula (A-6a) or (A-7a) or precursors thereof.
- In the definitions of symbols in this specification including the accompanying
- 30 claims unless otherwise specified the following definitions generally apply to the radicals in the formulae (I), (Ia) and (Ib) below:-
- 'halogen' means a fluorine, chlorine, bromine or iodine atom; and
- alkyl groups and moieties are straight or branched chain and contain from 1 to 6 carbon atoms.

Preferably A is a group of formula (A-1), (A-2), (A-3) or (A-4) (compounds of formula (A-1) are most preferred).

The benzoyl ring of the compounds of formula (I) is preferably 2,4-disubstituted, 2,3-disubstituted or 2,3,4-trisubstituted.

- 5 Preferably in formulae (A-4) to (A-7), the groups R^{14} , R^{15} , R^{16} , R^{100} , R^{101} , R^{102} , R^{14a} , R^{15a} , R^{16a} , R^{103} , R^{14b} , R^{15b} , R^{16b} , R^{104} , R^{105} and R^{108} are each hydrogen or lower alkyl (preferably hydrogen, methyl or ethyl); L (in A-7a) is NH; and Q is hydroxy or -S-phenyl.

- Compounds of formula (I) in which A is (A-1), (A-2) or (A-3); R is hydrogen or
10 $-CO_2R^3$ (in A-1 or A-2) wherein R^3 is a straight- or branched chain alkyl group containing up to three carbon atoms; and R^1 is cyclopropyl are preferred.

A further preferred class of compounds of formula (I) wherein A is (A-1) are those wherein:

R is hydrogen or $-CO_2Et$;

- 15 R^1 is cyclopropyl;
and two groups R^2 , on adjacent carbon atoms of the phenyl ring may, together with the carbon atoms to which they are attached, combine to form a 5 or 6 membered saturated or unsaturated heterocyclic ring which is fused to the 2,3 or 3,4 positions of the benzoyl ring; wherein the heterocyclic ring contains two hetero
20 atoms selected from sulphur and oxygen which are attached to the 2 and 3, or 3 and 4 positions of the benzoyl ring; and in which the 4-substituent of the benzoyl ring is halogen or $S(O)_pMe$, or the 2-substituent of the benzoyl ring is methyl, $S(O)_pMe$ or $-CH_2S(O)_qMe$ respectively; and optionally the heterocyclic ring may be substituted by one or more halogen atoms.

- 25 A further preferred class of compounds of formula (I) are those wherein A is (A-1); R is hydrogen or $-CO_2Et$; R^1 is cyclopropyl; R^2 is a halogen atom or a group selected from $-CF_3$, Me, Et, $-S(O)_pMe$, $-CH_2S(O)_qMe$ and optionally halogenated methoxy or ethoxy; and z is two or three.


- A further preferred class of compounds of formula (I) wherein A is (A-4) are those
30 wherein:

R^{14} , R^{15} , R^{16} , R^{100} , R^{101} and R^{102} are each hydrogen;

and two groups R^2 , on adjacent carbon atoms of the phenyl ring may, together with the carbon atoms to which they are attached, combine to form a 6 membered

saturated heterocyclic ring which is fused to the 2,3 or 3,4 positions of the benzoyl ring; wherein the heterocyclic ring contains a sulphur atom attached to the 4 position of the benzoyl ring, optionally in the form of a group -SO- or -SO₂-, and which ring is substituted by a 5- or 6- membered acetal thereof.

- 5 A more preferred class of compounds of formula (I) having the formula (Ia):



are those wherein:

R is hydrogen or -CO₂Et;


- 10 R¹¹⁷ is selected from -S(O)_pMe, Me, Et, a chlorine, bromine or fluorine atom, methoxy, ethoxy and -CH₂S(O)_qMe;

R¹¹⁸ is selected from a hydrogen atom, a chlorine, bromine or fluorine atom, methoxy, ethoxy and -S(O)_pMe;

- 15 R¹¹⁹ is selected from a hydrogen atom, a chlorine, bromine or fluorine atom, methoxy and CF₃; and

p and q each independently have the values zero, one or two.

An especially preferred class of compounds of formula (I) have the formula (Ib):




20

wherein R¹²⁰ is chlorine, bromine or trifluoromethyl; and


R is hydrogen or -CO₂Et.

Preferred diones are those in which a substituted phenyl ring as defined in formula (I); (Ia); or (Ib), is attached to a grouping;



Such diones in which the phenyl ring is substituted by two groups independently selected from halogen, alkyl, S(O)_palkyl (p = 0, 1 or 2) and haloalkyl are also preferred.

- 5 Preferred triones are those in which a substituted phenyl ring, as defined above, is attached to a grouping;



The following compounds of formula (I) are among the most preferred for use in the present invention:


- 10 5-cyclopropyl-4-[2-chloro-3-ethoxy-4-(ethylsulphonyl)benzoyl]isoxazole;
4-(4-chloro-2-methylsulphonylbenzoyl)-5-cyclopropylisoxazole;
5-cyclopropyl-4-(2-methylsulphonyl-4-trifluoromethylbenzoyl)isoxazole;
4-(4-bromo-2-methylsulphonylbenzoyl)-5-cyclopropylisoxazole;
5-cyclopropyl-4-[4-fluoro-3-methoxy-2-(methylsulphonyl)benzoyl]isoxazole;
15 4-(4-bromo-2-methylsulphonylmethylbenzoyl)-5-cyclopropylisoxazole;
ethyl 5-cyclopropyl-4-(2-methylsulphonyl-4-trifluoromethylbenzoyl)isoxazole-3-carboxylate;
2-[2-chloro-(4-methylsulphonyl)benzoyl]-1,3-cyclohexanedione;
2-[2-nitro-(4-methylsulphonyl)benzoyl]-1,3-cyclohexanedione;
20 2-(2,3-dihydro-5,8-dimethyl-1,1-dioxospiro[4H-1-benzothiine-4,2'-[1,3]dioxolan]-6-ylcarbonyl)cyclohexane-1,3-dione;
5-cyclopropyl-4-(2-methylsulphonyl-4-trifluoromethylbenzoyl)-3-methylthioisoxazole; and
2-cyano-3-cyclopropyl-1-(2-methylsulphonyl-4-trifluoromethylphenyl)propan-1,3-dione.
25

The most preferred compounds are 5-cyclopropyl-4-(2-methylsulphonyl-4-trifluoromethylbenzoyl)isoxazole and 2-[2-nitro-(4-methylsulphonyl)benzoyl]-1,3-cyclohexanedione.


Herbicidal benzoylisoxazole and/or dione compounds for use in this invention may be prepared by the methods described in the aforementioned patent publications, or by the application or adaptation of known methods used or described in the chemical literature.

- 5 It has been found that the aforementioned antidote compounds can be selected from a wide range of chemical substances. The preferred compositions of this invention may include one or more antidotes which are not suggested by earlier antidotes which have been proposed for use with the benzoylisoxazole and/or dione derivatives of formula (I). The compositions of the invention may include one
10 or more of the following antidotes:

a) compounds of the formulae (II) and (III),



(II)




(III)

20 wherein:


n' is an integer from zero to 5, preferably zero to 3;

T is a (C₁ or C₂)-alkanediyl chain which is unsubstituted or substituted by one or two (C₁-C₄)-alkyl radicals or by [(C₁-C₃)-alkoxy]carbonyl;


W is an unsubstituted or substituted divalent heterocyclic radical selected from the
25 group of the partially unsaturated or aromatic five-membered heterocyclic rings which have 1 to 3 hetero ring atoms of the N or O type, where the ring contains at least one N atom and not more than one O atom, preferably a radical selected from the group consisting of (W1) to (W4),




(W1)



(W2)



(W3)



(W4)

m' is zero or 1;

R^{17} and R^{19} are the same or different halogen, (C₁-C₄)-alkyl, (C₁-C₄)-alkoxy, nitro or (C₁-C₄)-haloalkyl;

R^{18} and R^{20} are the same or different OR²⁴, SR²⁴ or NR²⁴R²⁵ or a saturated or unsaturated 3- to 7-membered heterocycle having at least one N atom and up to 3 hetero atoms, preferably from the group selected from O and S, which is linked to the carbonyl group in (II) or (III) via the N atom and is unsubstituted or substituted by radicals selected from the group consisting of (C₁-C₄)-alkyl, (C₁-C₄)-alkoxy or optionally substituted phenyl, preferably a radical of the formula OR²⁴, NHR²⁵ or N(CH₃)₂, in particular of the formula OR²⁴;

R^{24} is hydrogen or an unsubstituted or substituted aliphatic hydrocarbon radical, preferably having a total of 1 to 18 C atoms;

R^{25} is hydrogen, (C₁-C₆)-alkyl, (C₁-C₆)-alkoxy or substituted or unsubstituted phenyl;

R^{26} is hydrogen, (C₁-C₈)-alkyl, (C₁-C₈)-haloalkyl, (C₁-C₄)-alkoxy-(C₁-C₄)-alkyl, (C₁-C₆)-hydroxyalkyl, (C₃-C₁₂)-cycloalkyl or tri-(C₁-C₄)-alkyl-silyl; and

R^{27} , R^{28} , R^{29} are the same or different hydrogen, (C₁-C₈)-alkyl, (C₁-C₈)-haloalkyl, (C₃-C₁₂)-cycloalkyl or substituted or unsubstituted phenyl;

b) one or more compounds selected from:

4,6-dichloro-2-phenylpyrimidine (fencloirim),

benzyl 2-chloro-4-trifluoromethyl-1,3-thiazole-5-carboxylate (flurazole),

N-(4-methylphenyl)-N'-(1-methyl-1-phenylethyl)urea (daimuron),

1-[4-(N-2-methoxybenzoylsulfamoyl)phenyl]-3-methylurea,

1-[4-(N-2-methoxybenzoylsulfamoyl)phenyl]-3,3-dimethylurea,

1-[4-(N-4,5-dimethylbenzoylsulfamoyl)phenyl]-3-methylurea,

1-[4-(N-naphthoylsulfamoyl)phenyl]-3,3-dimethylurea; or

(2,4-dichlorophenoxy)acetic acid (2,4-D),

(4-chlorophenoxy)acetic acid,

(R,S)-2-(4-chloro-o-tolyloxy)propionic acid (mecoprop),

4-(2,4-dichlorophenoxy)butyric acid (2,4-DB),

(4-chloro-o-tolyloxy)acetic acid (MCPA),

4-(4-chloro-o-tolyloxy)butyric acid,

4-(4-chlorophenoxy)butyric acid,


3,6-dichloro-2-methoxybenzoic acid (dicamba),

1-(ethoxycarbonyl)ethyl 3,6-dichloro-2-methoxybenzoate (lactidichlor)

and their salts and esters, preferably (C₁-C₈); or

- (±)-4-dichloroacetyl-3,4-dihydro-3-methyl-2*H*-1,4-benzoxazine (benoxacor),
N,N-diallyl-2,2-dichloroacetamide (dichlormid),
 (RS)-3-dichloroacetyl-5-(2-furyl)-2,2-dimethyloxazolidine (furilazole),
N-(4-chlorophenyl)maleimide (CMPI),
 5 4-hydroxy-1-methyl-3-(1-1*H*-tetrazol-5-ylmethanoyl)-1*H*-quinolin-2-one,
 (S)-1-(1- α -methylbenzyl)-3-*p*-tolylurea ((S)-MBU), and
 S-1-methyl-1-phenylethyl piperidine-1-carbothioate (dimepiperate).

c) N-acylsulfonamides of the formula (IV) or salts thereof:



(IV)

wherein:

- R^{30} is hydrogen; or R^{30} is a hydrocarbon radical, a hydrocarbon-oxy radical, a hydrocarbon-thio radical or a heterocyclyl radical which is preferably bonded via a carbon atom, which radicals are unsubstituted or substituted by one or more groups selected from halogen, cyano, nitro, amino, hydroxyl, carboxyl, formyl, carboxamide, sulfonamide and radicals of the formula Z^a-R^a , each hydrocarbon moiety preferably having 1 to 20 carbon atoms and a carbon-containing radical R^{30} inclusive of substituents preferably having 1 to 30 carbon atoms;
 15 R^{31} is hydrogen or (C₁-C₄)-alkyl, preferably hydrogen, or
 R^{30} and R^{31} together with the group of the formula -CO-N- are the residue of a 3- to 8-membered saturated or unsaturated ring;
 R^{32} is identical or different halogen, cyano, nitro, amino, hydroxyl, carboxyl, formyl, CONH₂, SO₂NH₂ or a radical of the formula Z^b-R^b ;
 25 R^{33} is hydrogen or (C₁-C₄)-alkyl, preferably H;
 R^{34} is the same or different halogen, cyano, nitro, amino, hydroxyl, carboxyl, CHO, CONH₂, SO₂NH₂ or a radical of the formula Z^c-R^c ;
 R^a is a hydrocarbon radical or a heterocyclyl radical, which radicals are unsubstituted or substituted by one or more groups which may be the same or
 30 different selected from halogen, cyano, nitro, amino, hydroxyl, mono- and di-[(C₁-C₄)-alkyl]amino, or an alkyl radical in which a plurality, preferably 2 or 3, non-adjacent CH₂ groups are in each case replaced by one oxygen atom;


R^b and R^c are the same or different hydrocarbon radical or heterocyclyl radical, which radicals are unsubstituted or substituted by one or more groups which may be the same or different selected from halogen, cyano, nitro, amino, hydroxyl, phosphoryl, halo-(C₁-C₄)-alkoxy, mono- and di-[(C₁-C₄)-alkyl]amino, or an alkyl radical in which a plurality, preferably 2 or 3; non-adjacent CH₂ groups are replaced in each case by one oxygen atom;

Z^a is a divalent group of the formula O, S, CO, CS, CO-O, CO-S, O-CO, S-CO, SO, SO₂, NR*, CO-NR*, NR*-CO, SO₂-NR* or NR*-SO₂, the bond given on the right-hand side of each of the divalent groups being the bond to the radical R^a , and the radicals R^* are each independently H, (C₁-C₄)-alkyl or halo-(C₁-C₄)-alkyl;

Z^b and Z^c are each independently a direct bond or a divalent group of the formula O, S, CO, CS, CO-O, CO-S, O-CO, S-CO, SO, SO₂, NR*, SO₂-NR*, NR*-SO₂, CO-NR* or NR*-CO, where, in asymmetrical divalent groups, the atom on the right-hand side is linked to the radical R^b or R^c and where the radicals R^* are each independently H, (C₁-C₄)-alkyl or halo-(C₁-C₄)-alkyl;

n is an integer from zero to 4, preferably zero, 1 or 2, in particular zero or 1, and t is an integer from zero to 5, preferably zero, 1, 2 or 3, in particular zero, 1 or 2;

d) acylsulfamoylbenzamides of the formula (V), or salts thereof:



(V)

wherein:

X^3 is CH or N;

R^{35} is hydrogen, or a heterocyclyl or hydrocarbon radical, which radicals are optionally substituted by one or more groups which may be the same or different selected from halogen, cyano, nitro, amino, hydroxyl, carboxyl, CHO, CONH₂, SO₂NH₂ and Z^d -R^d;

R^{36} is hydrogen, hydroxyl; or R^{36} is (C₁-C₆)-alkyl, (C₂-C₆)-alkenyl, (C₂-C₆)-alkynyl, (C₁-C₆)-alkoxy, (C₂-C₆)-alkenyloxy, which radicals are optionally substituted by one or more groups which may be the same or different selected from halogen, hydroxyl, (C₁-C₄)-alkyl, (C₁-C₄)-alkoxy and (C₁-C₄)-alkylthio, or

R^{35} and R^{36} together with the nitrogen atom to which they are attached are a 3- to 8-membered saturated or unsaturated ring;

R^{37} is halogen, cyano, nitro, amino, hydroxyl, carboxyl, CHO, CONH₂, SO₂NH₂ or Z^e-R^e;

5 R^{38} is hydrogen, (C₁-C₄)-alkyl, (C₂-C₄)-alkenyl or (C₂-C₄)-alkynyl;

R^{39} is halogen, cyano, nitro, amino, hydroxyl, carboxyl, phosphoryl, CHO, CONH₂, SO₂NH₂ or Z^f-R^f;

R^d is a (C₂-C₂₀)-alkyl radical whose carbon chain is interrupted by one or more oxygen atoms; or R^d is a heterocyclyl or a hydrocarbon radical, which radicals are optionally substituted by one or more groups which may be the same or different selected from halogen, cyano, nitro, amino, hydroxyl, mono- and di-[(C₁-C₄)-alkyl]amino;

10 R^e and R^f are the same or different (C₂-C₂₀)-alkyl radical which carbon chain is interrupted by one or more oxygen atoms; or R^e and R^f are the same or different heterocyclyl or hydrocarbon radical, which radicals are optionally substituted by one or more groups which may be the same or different selected from halogen, cyano, nitro, amino, hydroxyl, phosphoryl, (C₁-C₄)-haloalkoxy, mono- and di-[(C₁-C₄)-alkyl]amino;

20 Z^d is a divalent unit selected from the group consisting of O, S, CO, CS, C(O)O, C(O)S, SO, SO₂, NR*, C(O)NR* or SO₂NR*;


Z^e and Z^f are the same or different and are a direct bond or a divalent unit selected from O, S, CO, CS, C(O)O, C(O)S, SO, SO₂, NR*, SO₂NR* and C(O)NR*;

R* is hydrogen, (C₁-C₄)-alkyl or (C₁-C₄)-haloalkyl;

s is an integer from zero to 4, and

25 o is an integer from zero to 5 when X is CH, or is an integer from zero to 4 when X is N;

e) compounds of the formula (VI):



(VI)


wherein:

R^{40} is H, (C₁-C₄)-alkyl, (C₁-C₄)-alkyl substituted by (C₁-C₄)-alkyl-X⁴ or (C₁-C₄)-haloalkyl-X⁴, (C₁-C₄)-haloalkyl, NO₂, CN, -COO-R⁴³, NR₂⁴⁴, SO₂NR₂⁴⁵ or CONR₂⁴⁶;

R^{41} is H, halogen, (C₁-C₄)-alkyl, CF₃, (C₁-C₄)-alkoxy or (C₁-C₄)-haloalkoxy;

R^{42} is H, halogen or (C₁-C₄)-alkyl;

- 5 Q^1 , Q^2 , E and G are the same or different O, S, CR₂⁴⁷, CO, NR₂⁴⁸ or a group of the formula (VII):



(VII)

with the proviso that:

- 10 i) at least one of the groups Q^1 , Q^2 , E, G is a carbonyl group, that exactly one of these groups is a radical of the formula (VII) and that the group of the formula (VII) is adjacent to a carbonyl group, and
 ii) two adjacent groups Q^1 , Q^2 , E and G cannot simultaneously be oxygen;
- 15 R^9 is the same or different H or (C₁-C₈)-alkyl or the two radicals R^9 together are (C₂-C₆)-alkylene;
 V is Y³-R^h or NR₂⁴⁹;
 X^4 is O or S(O)_i;
 Y^3 is O or S;
- 20 R^h is H, (C₁-C₈)-alkyl, (C₁-C₈)-haloalkyl, (C₁-C₄)-alkoxy-(C₁-C₈)-alkyl, (C₃-C₆)-alkenyloxy-(C₁-C₈)-alkyl, or phenyl-(C₁-C₈)-alkyl, where the phenyl ring is optionally substituted by halogen, (C₁-C₄)-alkyl, CF₃, methoxy or methyl-S(O)_x; (C₃-C₆)-alkenyl, (C₃-C₆)-haloalkenyl, phenyl-(C₃-C₆)-alkenyl, (C₃-C₆)-alkynyl, phenyl-(C₃-C₆)-alkynyl, oxetanyl, furfuryl or tetrahydrofuryl;
- 25 R^{43} is H or (C₁-C₄)-alkyl;
 R^{44} is H, (C₁-C₄)-alkyl, (C₁-C₄)-alkylcarbonyl or the two radicals R^{44} together are (C₄-C₅)-alkylene;
- R^{45} and R^{46} are the same or different H, (C₁-C₄)-alkyl, or the two radicals R^{45} and/or R^{46} together are (C₄-C₅)-alkylene, where one CH₂ group can be replaced by O or S
- 30 or one or two CH₂ groups can be replaced by NRⁱ;
 R^i is H or (C₁-C₈)-alkyl;
 R^{47} is H, (C₁-C₈)-alkyl or the two radicals R^{47} together are (C₂-C₆)-alkylene;

R⁴⁸ is H, (C₁-C₈)-alkyl, substituted or unsubstituted phenyl, or benzyl which is unsubstituted or substituted on the phenyl ring;

R⁴⁹ is H, (C₁-C₈)-alkyl, phenyl, phenyl-(C₁-C₈)-alkyl, which phenyl rings can be substituted by F, Cl, Br, NO₂, CN, OCH₃, (C₁-C₄)-alkyl or CH₃SO₂; (C₁-C₄)-alkoxy-
 5 (C₁-C₈)-alkyl, (C₃-C₆)-alkenyl, (C₃-C₆)-alkynyl, (C₃-C₆)-cycloalkyl or two radicals R⁴⁹ together are (C₄-C₅)-alkylene, where one CH₂ group can be replaced by O or S or one or two CH₂ groups can be replaced by NR^k;

R^k is H or (C₁-C₄)-alkyl;

m'' is 0 or 1 and

10 l is 0, 1 or 2;

including stereoisomers and agriculturally acceptable salts.

Unless otherwise defined individually, the following definitions generally apply to the radicals in the formulae (II) to (VII).

15 The radicals alkyl, alkoxy, haloalkyl, haloalkoxy, alkylamino and alkylthio and the corresponding unsaturated and/or substituted radicals can be in each case straight-chain or branched in the carbon skeleton. Alkyl radicals, also the composite meanings such as alkoxy, haloalkyl and the like, preferably have 1 to 4 carbon atoms and are, for example, methyl, ethyl, n- or i-propyl or n-, i-, t- or 2-butyl.

20 Alkenyl and alkynyl radicals have the meanings of the unsaturated radicals which are possible which correspond to the alkyl radicals; alkenyl is, for example, allyl, 1-methylprop-2-en-1-yl, 2-methylprop-2-en-1-yl, but-2-en-1-yl, but-3-en-1-yl, 1-methylbut-3-en-1-yl and 1-methylbut-2-en-1-yl. Alkynyl is, for example, propargyl, but-2-yn-1-yl, but-3-yn-1-yl, 1-methylbut-3-yn-1-yl. "(C₁-C₄)-Alkyl" is the
 25 abbreviation for alkyl having 1 to 4 carbon atoms; the same applies analogously to other general definitions of radicals, where the range of the possible number of carbon atoms is indicated in brackets.

Cycloalkyl is, preferably, a cyclic alkyl radical having 3 to 8, preferably 3 to 7, especially preferably 3 to 6, carbon atoms, for example cyclopropyl, cyclobutyl,
 30 cyclopentyl and cyclohexyl. Cycloalkenyl and cycloalkynyl denote corresponding unsaturated compounds.

Halogen is fluorine, chlorine, bromine or iodine. Haloalkyl, haloalkenyl and haloalkynyl are alkyl, alkenyl or alkynyl which are partially or fully substituted by halogen, preferably by fluorine, chlorine and/or bromine, in particular by fluorine or
 35 chlorine, for example CF₃, CHF₂, CH₂F, CF₃CF₂, CH₂FCHCl, CCl₃, CHCl₂, CH₂CH₂Cl.

Haloalkoxy is, for example, OCF_3 , OCHF_2 , OCH_2F , $\text{CF}_3\text{CF}_2\text{O}$, OCH_2CF_3 and $\text{OCH}_2\text{CH}_2\text{Cl}$. This also applies analogously to other halogen-substituted radicals.

A hydrocarbon radical can be an aromatic or an aliphatic hydrocarbon radical, where an aliphatic hydrocarbon radical is generally a straight-chain or branched saturated or unsaturated hydrocarbon radical, preferably having 1 to 18, especially preferably 1 to 12, carbon atoms, for example alkyl, alkenyl or alkynyl.

Aliphatic hydrocarbon radical preferably means alkyl, alkenyl or alkynyl having up to 12 carbon atoms; the same applies analogously to an aliphatic hydrocarbon radical in a hydrocarbon-oxy radical.

Aryl is generally a mono-, bi- or polycyclic aromatic system having by preference 6-20 carbon atoms, preferably 6 to 14 carbon atoms, especially preferably 6 to 10 carbon atoms, for example phenyl, naphthyl, tetrahydronaphthyl, indenyl, indanyl, pentalenyl and fluorenyl, especially preferably phenyl.

Heterocyclic ring, heterocyclic radical or heterocyclyl is a mono-, bi- or polycyclic ring system which is saturated, unsaturated and/or aromatic and contains one or more, preferably 1 to 4, hetero atoms, preferably selected from the group consisting of N, S and O.

Preferred are saturated heterocycles having 3 to 7 ring atoms and one or two hetero atoms selected from the group consisting of N, O and S, their chalcogens not being adjacent. Especially preferred are monocyclic rings having 3 to 7 ring atoms and a hetero atom selected from the group consisting of N, O and S, and also morpholine, dioxolane, piperazine, imidazoline and oxazolidine. Very especially preferred saturated heterocycles are oxirane, pyrrolidone, morpholine and tetrahydrofuran.

Also preferred are partially unsaturated heterocycles having 5 to 7 ring atoms and one or two hetero atoms selected from the group consisting of N, O and S. Especially preferred are partially unsaturated heterocycles having 5 to 6 ring atoms and one hetero atom selected from the group consisting of N, O and S. Very especially preferred partially unsaturated heterocycles are pyrazoline, imidazoline and isoxazoline.

Equally preferred is heteroaryl, for example mono- or bicyclic aromatic heterocycles having 5 to 6 ring atoms which contain one to four hetero atoms selected from the group consisting of N, O, S, the chalcogens not being adjacent. Especially preferred are monocyclic aromatic heterocycles having 5 to 6 ring atoms which contains a hetero atom selected from the group consisting of N, O and S, and also

pyrimidine, pyrazine, pyridazine, oxazole, thiazole, thiadiazole, oxadiazole, pyrazole, triazole and isoxazole. Very especially preferred are pyrazole, thiazole, triazole and furan.

Substituted radicals, such as substituted hydrocarbon radicals, for example
5 substituted alkyl, alkenyl, alkynyl, aryl such as phenyl and arylalkyl such as benzyl, or substituted heterocyclyl, are a substituted radical which is derived from an unsubstituted skeleton, the substituents being, by preference, one or more, by preference 1, 2 or 3, in the case of Cl and F also up to the maximum possible number of, substituents selected from the group consisting of halogen, alkoxy,
10 haloalkoxy, alkylthio, hydroxyl, amino, nitro, carboxyl, cyano, azido, alkoxy-carbonyl, alkylcarbonyl, formyl, carbamoyl, mono- and dialkylaminocarbonyl, substituted amino such as acylamino, mono- and dialkylamino and alkylsulfynyl, haloalkylsulfynyl, alkylsulfonyl, haloalkylsulfonyl and, in the case of cyclic radicals, also alkyl and haloalkyl and the unsaturated aliphatic substituents which
15 correspond to the abovementioned saturated hydrocarbon-containing substituents, preferably alkenyl, alkynyl, alkenyloxy and alkynyloxy. In the case of radicals having carbon atoms, those having 1 to 4 carbon atoms, in particular 1 or 2 carbon atoms, are preferred. As a rule, preferred substituents are those selected from the group consisting of halogen, for example fluorine or chlorine, (C₁-C₄)-alkyl,
20 preferably methyl or ethyl, (C₁-C₄)-haloalkyl, preferably trifluoromethyl, (C₁-C₄)-alkoxy, preferably methoxy or ethoxy, (C₁-C₄)-haloalkoxy, nitro and cyano. Especially preferred in this context are the substituents methyl, methoxy and chlorine.

Mono- or disubstituted amino is a chemically stable radical selected from the group
25 of the substituted amino radicals which are N-substituted by, for example, one or two identical or different radicals selected from the group consisting of alkyl, alkoxy, acyl and aryl; preferably monoalkylamino, dialkylamino, acylamino, arylamino, N-alkyl-N-aryl-amino and N-heterocycles. Preferred in this context are alkyl radicals having 1 to 4 carbon atoms. By preference, aryl is phenyl. By
30 preference, substituted aryl is substituted phenyl. The definition given further below applies to acyl, preferably (C₁-C₄)-alkanoyl. This also applies analogously to substituted hydroxylamino or hydrazino.

By preference, optionally substituted phenyl is phenyl which is unsubstituted or
35 mono- or polysubstituted, preferably up to trisubstituted, in the case of halogen such as Cl and F also up to pentasubstituted, by identical or different radicals

selected from the group consisting of halogen, (C₁-C₄)-alkyl, (C₁-C₄)-alkoxy, (C₁-C₄)-haloalkyl, (C₁-C₄)-haloalkoxy and nitro, for example o-, m- and p-tolyl, dimethylphenyls, 2-, 3- and 4-chlorophenyl, 2-, 3- and 4-trifluoro- and -trichlorophenyl, 2,4-, 3,5-, 2,5- and 2,3-dichlorophenyl, o-, m- and p-methoxyphenyl.

An acyl radical is the radical of an organic acid having by preference up to 6 carbon atoms, for example the radical of a carboxylic acid and radicals of acids derived therefrom, such as thiocarboxylic acid, optionally N-substituted iminocarboxylic acids, or the radical of carbonic monoesters, optionally N-substituted carbamic acids, sulfonic acid, sulfinic acids, phosphonic acids, phosphinic acids. Acyl is, for example, formyl, alkylcarbonyl such as (C₁-C₄-alkyl)-carbonyl, phenylcarbonyl, it being possible for the phenyl ring to be substituted, for example as indicated above for phenyl, or alkyloxycarbonyl, phenyloxycarbonyl, benzyloxycarbonyl, alkylsulfonyl, alkylsulfonyl or N-alkyl-1-iminoalkyl.

Formulae (II) to (VII) are also understood to include all stereoisomers and mixtures thereof. The stereoisomers include enantiomers, diastereomers and Z- and E-isomers.

Preferred herbicide/antidote combinations are those which comprise antidotes of the formula (II) and/or (III) where the symbols and indices have the following meanings:

R²⁴ is hydrogen, (C₁-C₁₈)-alkyl, (C₃-C₁₂)-cycloalkyl, (C₂-C₈)-alkenyl and (C₂-C₁₈)-alkynyl, where the carbon-containing groups can be substituted by one or more preferably up to three, radicals R⁵⁰;

R⁵⁰ is the same or different halogen, hydroxyl, (C₁-C₈)-alkoxy, (C₁-C₈)-alkylthio, (C₂-C₈)-alkenylthio, (C₂-C₈)-alkynylthio, (C₂-C₈)-alkenyloxy, (C₂-C₈)-alkynyloxy, (C₃-C₇)-cycloalkyl, (C₃-C₇)-cycloalkoxy, cyano, mono- and di-(C₁-C₄)-alkylamino, carboxyl, (C₁-C₈)-alkoxycarbonyl, (C₂-C₈)-alkenyloxycarbonyl, (C₁-C₈)-alkylthiocarbonyl, (C₂-C₈)-alkynyloxycarbonyl, (C₁-C₈)-alkylcarbonyl, (C₂-C₈)-alkenylcarbonyl, (C₂-C₈)-alkynylcarbonyl, 1-(hydroxyimino)-(C₁-C₆)-alkyl, 1-[(C₁-C₄)-alkylimino]-(C₁-C₄)-alkyl, 1-[(C₁-C₄)-alkoxyimino]-(C₁-C₆)-alkyl, (C₁-C₈)-alkylcarbonylamino, (C₂-C₈)-alkenylcarbonylamino, (C₂-C₈)-alkynylcarbonylamino, aminocarbonyl, (C₁-C₈)-alkylaminocarbonyl, di-(C₁-C₆)-alkylaminocarbonyl, (C₂-C₆)-alkenylaminocarbonyl, (C₂-C₆)-alkynylaminocarbonyl, (C₁-C₈)-alkoxycarbonylamino, (C₁-C₈)-

- alkylaminocarbonylamino, (C₁-C₆)-alkylcarbonyloxy which is unsubstituted or substituted by R⁵¹, or is (C₂-C₆)-alkenylcarbonyloxy, (C₂-C₆)-alkynylcarbonyloxy, (C₁-C₈)-alkylsulfonyl, phenyl, phenyl-(C₁-C₆)-alkoxy, phenyl-(C₁-C₆)-alkoxycarbonyl, phenoxy, phenoxy-(C₁-C₆)-alkoxy, phenoxy-(C₁-C₆)-alkoxycarbonyl,
- 5 phenylcarbonyloxy, phenylcarbonylamino, phenyl-(C₁-C₆)-alkylcarbonylamino, it being possible for the last-mentioned 9 radicals to be unsubstituted or mono- or polysubstituted, preferably up to trisubstituted, by radicals R⁵²; SiR'₃, OSiR'₃, R'₃Si- (C₁-C₈)-alkoxy, CO-O-NR'₂, O-N=CR'₂, N=CR'₂, O-N R'₂, NR'₂, CH(OR')₂, O-(CH₂)_q-CH(OR')₂, CR'''(OR')₂, O-(CH₂)_wCR'''(OR')₂ or by R''O-CHR'''CHCOR''-(C₁-C₆)-
- 10 alkoxy,
- R⁵¹ is the same or different halogen, nitro, (C₁-C₄)-alkoxy and phenyl which is unsubstituted or substituted by one or more, preferably up to three, radicals R⁵²; R⁵² is the same or different halogen, (C₁-C₄)-alkyl, (C₁-C₄)-alkoxy, (C₁-C₄)-haloalkyl, (C₁-C₄)-haloalkoxy or nitro;
- 15 R' is the same or different hydrogen, (C₁-C₄)-alkyl, phenyl which is unsubstituted or substituted by one or more, preferably up to three, radicals R⁵², or two radicals R' together form a (C₂-C₆)-alkanediyl chain;
- R'' is the same or different (C₁-C₄)-alkyl or two radicals R'' together form a (C₂-C₆)-alkanediyl chain;
- 20 R''' is hydrogen or (C₁-C₄)-alkyl; and
- w is zero, 1, 2, 3, 4, 5 or 6.

Especially preferred are herbicide/antidote combinations according to the invention which comprise antidotes of the formula (II) and/or (III) where the symbols and

25 indices have the following meanings:

- R²⁴ is hydrogen, (C₁-C₈)-alkyl or (C₃-C₇)-cycloalkyl, the above carbon-containing radicals being unsubstituted or mono- or polysubstituted by halogen or mono- or disubstituted, by preference monosubstituted, by radicals R⁵⁰,
- R⁵⁰ is the same or different hydroxyl, (C₁-C₄)-alkoxy, carboxyl, (C₁-C₄)-
- 30 alkoxycarbonyl, (C₂-C₆)-alkenyloxy, (C₂-C₆)-alkynyloxy, 1-(hydroxyimino)-(C₁-C₄)-alkyl, 1-[(C₁-C₄)-alkylimino]-(C₁-C₄)-alkyl and 1-[(C₁-C₄)-alkoxyimino]-(C₁-C₄)-alkyl; SiR'₃, O-N=CR'₂, N=CR'₂, NR'₂ and ONR'₃ where R' is identical or different hydrogen, (C₁-C₄)-alkyl or, as a pair, a (C₄-C₅)-alkanediyl chain, R²⁷, R²⁸ and R²⁹ are the same or different hydrogen, (C₁-C₈)-alkyl, (C₁-C₆)-haloalkyl,
- 35 (C₃-C₇)-cycloalkyl or phenyl which is unsubstituted or substituted by one or more

groups selected from halogen, cyano, nitro, amino, mono- and di-[(C₁-C₄)-alkyl]-amino, (C₁-C₄)-alkyl, (C₁-C₄)-haloalkyl, (C₁-C₄)-alkoxy, (C₁-C₄)-haloalkoxy, (C₁-C₄)-alkylthio and (C₁-C₄)-alkylsulfonyl;

R²⁶ is hydrogen, (C₁-C₈)-alkyl, (C₁-C₈)-haloalkyl, (C₁-C₄-alkoxy)-(C₁-C₄)-alkyl, (C₁-C₆)-hydroxyalkyl, (C₃-C₇)-cycloalkyl or tri-(C₁-C₄)-alkylsilyl,

R¹⁷ and R¹⁹ are the same or different halogen, methyl, ethyl, methoxy, ethoxy, (C₁ or C₂)-haloalkyl, by preference hydrogen, halogen or (C₁ or C₂)-haloalkyl.

Very especially preferred are antidotes in which the symbols and indices in formula (II) have the following meanings:

R¹⁷ is halogen, nitro or (C₁-C₄)-haloalkyl;

n' is zero, 1, 2 or 3;

R¹⁸ is a radical of the formula OR²⁴,

R²⁴ is hydrogen, (C₁-C₈)-alkyl or (C₃-C₇)-cycloalkyl, where the above carbon-

containing radicals are unsubstituted or mono- or polysubstituted, by preference up to trisubstituted, by the same or different halogen radicals, or up to disubstituted, by preference monosubstituted, by identical or different radicals selected from the group consisting of hydroxyl, (C₁-C₄)-alkoxy, (C₁-C₄)-alkoxycarbonyl, (C₂-C₆)-alkenyloxycarbonyl, (C₂-C₆)-alkynyloxycarbonyl,

1-(hydroxyimino)-(C₁-C₄)-alkyl, 1-[(C₁-C₄)-alkylimino]-(C₁-C₄)-alkyl, 1-[(C₁-C₄)-alkoxyimino]-(C₁-C₄)-alkyl and radicals of the formulae SiR'₃, O-N=CR'₂, N=CR'₂, NR'₂ and O-NR'₂, where the radicals R' in the abovementioned formulae are identical or different hydrogen, (C₁-C₄)-alkyl or, as a pair, are (C₄ or C₅)-alkanediyl;

R²⁷, R²⁸ and R²⁹ are the same or different hydrogen, (C₁-C₈)-alkyl, (C₁-C₆)-haloalkyl, (C₃-C₇)-cycloalkyl or phenyl which is unsubstituted or substituted by one or more radicals selected from halogen, (C₁-C₄)-alkyl, (C₁-C₄)-alkoxy, nitro, (C₁-C₄)-haloalkyl and (C₁-C₄)-haloalkoxy, and

R²⁶ is hydrogen, (C₁-C₈)-alkyl, (C₁-C₈)-haloalkyl, (C₁-C₄)-alkoxy-(C₁-C₄)-alkyl, (C₁-C₆)-hydroxyalkyl, (C₃-C₇)-cycloalkyl or tri-(C₁-C₄)-alkylsilyl.

Very especially preferred are also antidotes of the formula (III) where the symbols and indices have the following meanings:

R¹⁹ is halogen or (C₁-C₄)-haloalkyl;

n' is zero, 1, 2 or 3, where (R¹⁹)_{n'} is, by preference, 5-Cl;

R²⁰ is a radical of the formula OR²⁴;

T is CH₂ or CH(COO-(C₁-C₃)-alkyl), and

R²⁴ is hydrogen, (C₁-C₈)-alkyl, (C₁-C₈)-haloalkyl or (C₁-C₄)-alkoxy-(C₁-C₄)-alkyl, by preference (C₁-C₈)-alkyl.

- 5 Especially preferred are antidotes of the formula (II) where the symbols and indices have the following meanings:

W is (W1);

R¹⁷ is halogen or (C₁-C₂)-haloalkyl;

n' is zero, 1, 2 or 3, where (R¹⁷)_n is by preference 2,4-Cl₂;

- 10 R¹⁸ is a radical of the formula OR²⁴;

R²⁴ is hydrogen, (C₁-C₈)-alkyl, (C₁-C₄)-haloalkyl, (C₁-C₄)-hydroxyalkyl, (C₃-C₇)-cycloalkyl, (C₁-C₄)-alkoxy-(C₁-C₄)-alkyl or tri-(C₁-C₂)-alkylsilyl, by preference (C₁-C₄)-alkyl;

R²⁷ is hydrogen, (C₁-C₈)-alkyl, (C₁-C₄)-haloalkyl or (C₃-C₇)-cycloalkyl, by preference hydrogen or (C₁-C₄)-alkyl, and

- 15 R²⁶ is hydrogen, (C₁-C₈)-alkyl, (C₁-C₄)-haloalkyl, (C₁-C₄)-hydroxyalkyl, (C₃-C₇)-cycloalkyl, (C₁-C₄)-alkoxy-(C₁-C₄)-alkyl or tri-(C₁-C₂)-alkylsilyl, by preference hydrogen or (C₁-C₄)-alkyl.

- Also especially preferred are herbicidal compositions comprising an antidote of the
20 formula (II) where the symbols and indices have the following meanings:

W is (W2);

R¹⁷ is halogen or (C₁-C₂)-haloalkyl;

n' is zero, 1, 2 or 3, where (R¹⁷)_n is by preference 2,4-Cl₂;

R¹⁸ is a radical of the formula OR²⁴;

- 25 R²⁴ is hydrogen, (C₁-C₈)-alkyl, (C₁-C₄)-haloalkyl, (C₁-C₄)-hydroxyalkyl, (C₃-C₇)-cycloalkyl, (C₁-C₄)-alkoxy-(C₁-C₄)-alkyl or tri-(C₁-C₂)-alkyl-silyl, by preference (C₁-C₄)-alkyl, and

R²⁷ is hydrogen, (C₁-C₈)-alkyl, (C₁-C₄)-haloalkyl, (C₃-C₇)-cycloalkyl or phenyl, by preference hydrogen or (C₁-C₄)-alkyl.

30

Also especially preferred are antidotes of the formula (II) where the symbols and indices have the following meanings:

W is (W3);

R¹⁷ is halogen or (C₁-C₂)-haloalkyl;

- 35 n' is zero, 1, 2 or 3, where (R¹⁷)_n is by preference 2,4-Cl₂;

R^{18} is a radical of the formula OR^{24} ;

R^{24} is hydrogen, (C_1-C_8) -alkyl, (C_1-C_4) -haloalkyl, (C_1-C_4) -hydroxyalkyl, (C_3-C_7) -cycloalkyl, (C_1-C_4) -alkoxy- (C_1-C_4) -alkyl or tri- (C_1-C_2) -alkylsilyl, by preference (C_1-C_4) -alkyl, and

5 R^{28} is (C_1-C_8) -alkyl or (C_1-C_4) -haloalkyl, by preference C_1 -haloalkyl.

Also especially preferred are antidotes of the formula (II) where the symbols and indices have the following meaning:

W is (W4);

10 R^{17} is halogen, nitro, (C_1-C_4) -alkyl, (C_1-C_2) -haloalkyl, by preference CF_3 , or (C_1-C_4) -alkoxy;

n' is 0, 1, 2 or 3;

m' is 0 or 1;

R^{18} is a radical of the formula OR^{24} ;

15 R^{24} is hydrogen, (C_1-C_4) -alkyl, carboxy- (C_1-C_4) -alkyl, (C_1-C_4) -alkoxycarbonyl- (C_1-C_4) -alkyl, by preference (C_1-C_4) -alkoxy- $CO-CH_2-$, (C_1-C_4) -alkoxy- $CO-C(CH_3)(H)-$, $HO-CO-CH_2-$ or $HO-CO-C(CH_3)(H)-$, and

R^{29} is hydrogen, (C_1-C_4) -alkyl, (C_1-C_4) -haloalkyl, (C_3-C_7) -cycloalkyl or phenyl which is unsubstituted or substituted by one or more groups selected from halogen, $(C_1-$

20 $C_4)$ -alkyl, (C_1-C_4) -haloalkyl, nitro, cyano and (C_1-C_4) -alkoxy.

The following groups of compounds are especially suitable for use as antidotes for the herbicidally active substances of the formula (I):

a) compounds of the dichlorophenylpyrazoline-3-carboxylic acid type (i.e. of the formula (II), where $W = (W1)$ and $(R^{17})_{n'} = 2,4-Cl_2$), by preference compounds such as ethyl 1-(2,4-dichlorophenyl)-5-(ethoxycarbonyl)-5-methyl-2-pyrazoline-3-carboxylate (II-1), and related compounds as described in WO-A 91/07874;

b) dichlorophenylpyrazolecarboxylic acid derivatives (i.e. of the formula (II), where $W = (W2)$ and $(R^{17})_{n'} = 2,4-Cl_2$), by preference compounds such as ethyl 1-(2,4-dichlorophenyl)-5-methyl-pyrazole-3-carboxylate (II-2), ethyl 1-(2,4-dichlorophenyl)-5-isopropylpyrazole-3-carboxylate (II-3), ethyl 1-(2,4-dichlorophenyl)-5-(1,1-dimethylethyl)pyrazole-3-carboxylate (II-4), ethyl 1-(2,4-dichlorophenyl)-5-phenylpyrazole-3-carboxylate (II-5) and related compounds as described in EP-A-O 333 131 and EP-A-O 269 806.

- c) Compounds of the triazolecarboxylic type (i.e. of the formula (II), where $W = (W3)$ and $(R^{17})_n = 2,4-Cl_2$), by preference compounds such as fenchlorazole ethyl, i.e. ethyl 1-(2,4-dichlorophenyl)-5-trichloromethyl-(1H)-1,2,4-triazole-3-carboxylate (II-6); fenchlorazole, i.e. 1-(2,4-dichlorophenyl)-5-trichloromethyl-(1H)-1,2,4-triazole-3-carboxylic acid, and related compounds (see EP-A-O 174 562 and EP-A-O 346 620);
- d) Compounds of the 5-benzyl- or 5-phenyl-2-isoxazoline-3-carboxylic acid type or the 5,5-diphenyl-2-isoxazoline-3-carboxylic acid type (where $W = (W4)$), by preference compounds such as ethyl 5-(2,4-dichlorobenzyl)-2-isoxazoline-3-carboxylate (II-7) or ethyl 5-phenyl-2-isoxazoline-3-carboxylate (II-8) and related compounds as they are described in WO-A- 91/08202, or ethyl 5,5-diphenyl-2-isoxazoline-3-carboxylate (II-9), or 5,5-diphenyl-2-isoxazoline-3-carboxylic acid, or n-propyl 5,5-diphenyl-2-isoxazoline-carboxylate (II-10) or ethyl 5-(4-fluorophenyl)-5-phenyl-2-isoxazoline-3-carboxylate (II-11), as described in WO-A-95/07897.
- e) Compounds of the 8-quinolinoxyacetic acid type, for example those of the formula (III) where $(R^{19})_n = 5-Cl$, $R^{20} = OR^{24}$ and $T = CH_2$, by preference the compounds
- 1-methyl (5-chloro-8-quinolinoxy)acetate (III-1),
1,3-dimethyl-but-1-yl (5-chloro-8-quinolinoxy)acetate (III-2),
4-allyloxybutyl (5-chloro-8-quinolinoxy)acetate (III-3),
1-allyloxyprop-2-yl (5-chloro-8-quinolinoxy)acetate (III-4),
ethyl (5-chloro-8-quinolinoxy)acetate (III-5),
methyl (5-chloro-8-quinolinoxy)acetate (III-6),
allyl (5-chloro-8-quinolinoxy)acetate (III-7),
2-(2-propylideneiminoxy)-1-ethyl (5-chloro-8-quinolinoxy)acetate (III-8),
2-oxo-prop-1-yl (5-chloro-8-quinolinoxy)acetate (III-9) and related compounds as they are described in EP-A-O 860 750, EP-A-O 094 349 and EP-A-O 191 736 or EP-A-O 492 366.
- f) Compounds of the (5-chloro-8-quinolinoxy)malonic acid type, i.e. of the formula (III) where $(R^{19})_n = 5-Cl$, $R^{20} = OR^{24}$, $T = -CH(COO-alkyl)-$, by preference the compounds diethyl (5-chloro-8-quinolinoxy)malonate, diallyl (5-chloro-8-

quinolinoxy)malonate, methyl ethyl (5-chloro-8-quinolinoxy)malonate and related compounds as they are described in EP-A-0 582 198.

g) Compounds selected from:

- 5 N,N-diallyl-2,2-dichloroacetamide (dichlormid, from US-A 4,137,070),
4-dichloroacetyl-3,4-dihydro-3-methyl-2H-1,4-benzoxazine (benoxacor, from EP 0 149 974),
3-dichloroacetyl-5-(2-furanyl)-2,2-dimethyloxazolidine (furilazole, MON 13900),
4,6-dichloro-2-phenylpyrimidine (fenclorim),
10 benzyl 2-chloro-4-trifluoromethyl-1,3-thiazole-5-carboxylate (flurazole),
N-(4-methylphenyl)-N'-(1-methyl-1-phenylethyl)urea (daimuron),
1-[4-(N-2-methoxybenzoylsulfamoyl)phenyl]-3-methylurea,
1-[4-(N-2-methoxybenzoylsulfamoyl)phenyl]-3,3-dimethylurea,
1-[4-(N-4,5-dimethylbenzoylsulfamoyl)phenyl]-3-methylurea,
15 1-[4-(N-naphthoylsulfamoyl)phenyl]-3,3-dimethylurea,
(2,4-dichlorophenoxy)acetic acid (2,4-D); or
(4-chlorophenoxy)acetic acid,
(R,S)-2-(4-chloro-o-tolyloxy)propionic acid (mecoprop),
4-(2,4-dichlorophenoxy)butyric acid (2,4-DB),
20 (4-chloro-o-tolyloxy)acetic acid (MCPA),
4-(4-chloro-o-tolyloxy)butyric acid,
4-(4-chlorophenoxy)butyric acid,
3,6-dichloro-2-methoxybenzoic acid (dicamba),
1-(ethoxycarbonyl)ethyl 3,6-dichloro-2-methoxybenzoate (lactidichlor),
25 and their salts and esters, by preference (C₁-C₈); or
N-(4-chlorophenyl)maleimide (CMPI),
4-hydroxy-1-methyl-3-(1-1H-tetrazol-5-ylmethanoyl)-1H-quinolin-2-one,
(S)-1-(1-alpha-methylbenzyl)-3-p-tolylurea ((S)-MBU), and
S-1-methyl-1-phenylethyl piperidine-1-carbothioate (dimepiperate).

30

Furthermore preferred as antidotes are compounds of the formula (IV) or salts thereof in which:

R³⁰ is hydrogen; or R³⁰ is (C₁-C₆)-alkyl, (C₃-C₆)-cycloalkyl, furanyl or thienyl, which radicals are unsubstituted or substituted by one or more groups selected from

halogen, (C₁-C₄)-alkoxy, halo-(C₁-C₆)-alkoxy and (C₁-C₄)-alkylthio and, in the case of cyclic radicals, also (C₁-C₄)-alkyl and (C₁-C₄)-haloalkyl,

R³¹ is hydrogen,

R³² is halogen, halo-(C₁-C₄)-alkyl, halo-(C₁-C₄)-alkoxy, (C₁-C₄)-alkyl, (C₁-C₄)-alkoxy,

5 (C₁-C₄)-alkylsulfonyl, (C₁-C₄)-alkoxycarbonyl or (C₁-C₄)-alkylcarbonyl, by preference halogen, (C₁-C₄)-haloalkyl such as trifluoromethyl, (C₁-C₄)-alkoxy, halo-(C₁-C₄)-alkoxy, (C₁-C₄)-alkoxycarbonyl or (C₁-C₄)-alkylsulfonyl,

R³³ is hydrogen,

R³⁴ is halogen, (C₁-C₄)-alkyl, halo-(C₁-C₄)-alkyl, halo-(C₁-C₄)-alkoxy, (C₃-C₆)-

10 cycloalkyl, phenyl, (C₁-C₄)-alkoxy, cyano, (C₁-C₄)-alkylthio, (C₁-C₄)-alkylsulfonyl, (C₁-C₄)-alkoxycarbonyl or (C₁-C₄)-alkylcarbonyl,

by preference halogen, (C₁-C₄)-alkyl, (C₁-C₄)-haloalkyl such as trifluoromethyl, halo-(C₁-C₄)-alkoxy, (C₁-C₄)-alkoxy or (C₁-C₄)-alkylthio,

n is zero, 1 or 2 and

15 t is 1 or 2.

Furthermore preferred are antidotes of the formula (V) in which

X³ is CH;

R³⁵ is hydrogen; or R³⁵ is (C₁-C₆)-alkyl, (C₃-C₆)-cycloalkyl, (C₂-C₆)-alkenyl, (C₅-C₆)-

20 cycloalkenyl, phenyl or 3- to 6-membered heterocyclyl having up to three hetero atoms selected from the group consisting of nitrogen, oxygen and sulfur, which radicals are optionally substituted by one or more groups which may be the same or different selected from halogen, (C₁-C₆)-alkoxy, (C₁-C₆)-haloalkoxy, (C₁-C₂)-alkylsulfonyl, (C₁-C₂)-alkylsulfonyl, (C₃-C₆)-cycloalkyl, (C₁-C₄)-alkoxycarbonyl, (C₁-C₄)-alkylcarbonyl and phenyl and in the case of cyclic radicals, also (C₁-C₄)-alkyl and (C₁-C₄)-haloalkyl;

R³⁶ is hydrogen; or R³⁶ is (C₁-C₆)-alkyl, (C₂-C₆)-alkenyl, (C₂-C₆)-alkynyl, which radicals are optionally substituted by one or more groups which may be the same or different selected from halogen, hydroxyl, (C₁-C₄)-alkyl, (C₁-C₄)-alkoxy and (C₁-C₄)-alkylthio;

R³⁷ is halogen, (C₁-C₄)-haloalkyl, (C₁-C₄)-haloalkoxy, nitro, (C₁-C₄)-alkyl, (C₁-C₄)-alkoxy, (C₁-C₄)-alkylsulfonyl, (C₁-C₄)-alkoxycarbonyl or (C₁-C₄)-alkylcarbonyl;

R³⁸ is hydrogen;

R³⁹ is halogen, nitro, (C₁-C₄)-alkyl, (C₁-C₄)-haloalkyl, (C₁-C₄)-haloalkoxy, (C₃-C₆)-cycloalkyl, phenyl, (C₁-C₄)-alkoxy, cyano, (C₁-C₄)-alkylthio, (C₁-C₄)-alkylsulfinyl, (C₁-C₄)-alkylsulfonyl, (C₁-C₄)-alkoxycarbonyl or (C₁-C₄)-alkylcarbonyl;

s is zero, 1 or 2, and

5 o is 1 or 2.

The following sub-groups are particularly preferred amongst the antidotes of the formula (VI):

- compounds in which R⁴⁸ and R⁴⁹ are H, (C₁-C₈)-alkyl, phenyl, phenyl-(C₁-C₈)-alkyl, (C₁-C₄)-alkoxy-(C₁-C₈)-alkyl, (C₃-C₆)-alkenyl or (C₃-C₆)-alkynyl, it being possible for phenyl rings to be substituted by F, Cl, Br, NO₂, CN, OCH₃, (C₁-C₄)-alkyl or CH₃-SO₂;
- compounds in which R⁹ is H;
- compounds in which V is Y-R^h;
- 15 - compounds in which E is oxygen;
- compounds in which Q¹ is CR₂⁴⁷;
- compounds in which R⁴⁷ is hydrogen;
- compounds where m'' = 1 and E is oxygen or sulfur;
- compounds in which m'' = 0;
- 20 - compounds in which R⁴⁰, R⁴¹, R⁴², R⁴³ and R⁴⁴ are in each case hydrogen, E is oxygen, Q¹ is CR₂⁴⁷, V is Y-R^h and m'' = 1, in particular those where R⁴⁷ is H, R^b is CH₃ and Y is oxygen;
- compounds in which Q¹ is CR₂⁴⁷ and m'' is zero, in particular those in which R⁴⁴ and R⁴⁷ are hydrogen and V is Y-R^h, where R^h is by preference methyl and Y is by preference oxygen.

The most especially preferred antidotes are flurazole, which is benzyl 2-chloro-4-trifluoromethyl-1,3-thiazole-5-carboxylate; fenchlorazole-ethyl, which is ethyl 1-(2,4-dichlorophenyl)-5-trichloromethyl-1*H*-1,2,4-triazole-3-carboxylate; fenchlorazole, which is 1-(2,4-dichlorophenyl)-5-trichloromethyl-1*H*-1,2,4-triazole-3-carboxylic acid; benoxacor, which is (±)-4-dichloroacetyl-3,4-dihydro-3-methyl-2*H*-1,4-benzoxazine; dichlormid, which is *N,N*-diallyl-2,2-dichloroacetamide; fencloir, which is 4,6-dichloro-2-phenylpyrimidine; furilazole, which is (*RS*)-3-dichloroacetyl-5-(2-furyl)-2,2-dimethyloxazolidine; mefenpyr-diethyl, which is diethyl (*RS*)-1-(2,4-dichlorophenyl)-5-methyl-2-pyrazoline-3,5-dicarboxylate; CMPI, which is *N*-(4-chlorophenyl)maleimide; 4-hydroxy-1-methyl-3-(1-1*H*-tetrazol-5-

ylmethanoyl)-1*H*-quinolin-2-one; daimuron, which is 1-(1-methyl-1-phenylethyl)-3-*p*-tolylurea; (S)-MBU, which is (S)-1-(1- α -methylbenzyl)-3-*p*-tolylurea; dimepiperate, which is S-1-methyl-1-phenylethyl piperidine-1-carbothioate; 5,5-diphenylisoxazoline-3-carboxylic acid; and ethyl 5,5-diphenylisoxazoline-3-carboxylate (most preferably fenclorazole; CMPI; 4-hydroxy-1-methyl-3-(1-1*H*-tetrazol-5-ylmethanoyl)-1*H*-quinolin-2-one; (S)-MBU and dimepiperate).

The mixtures of the invention may be used to obtain selective weed control with low crop injury in various crop plants such as maize, soybean, cotton, canola, sugar beet, potatoes, wheat, tobacco, rice and oil seed rape. Preferred crops include maize, sugar beet, cotton and canola. Particularly preferred crop species are maize and soybean, especially maize.

Effective weed control coupled with low crop injury is a result of treatment of a plant locus with a combination of a herbicidal benzoylisoxazole and/or dione derivative and an antidote compound in accordance with the method of the present invention. By application to the 'plant locus' is meant application, for example to the plant growing medium, such as soil, as well as to the seeds, emerging seedlings, roots, stems, leaves or other plant parts.

The phrase 'combination of a herbicidal isoxazole and/or dione derivative and an antidote compound' includes various methods of treatment. For example, the soil of a plant locus may be treated with a "tank-mix" composition containing a mixture of the herbicide and the antidote which is "in combination" , or the soil may be treated with the herbicide and antidote compounds separately so that the "combination" may be made on, or in the soil. After such treatments of the soil with a mixture of herbicide and antidote or by separate or sequential application of the herbicide and the antidote to the soil, the herbicide and antidote may be mixed into or incorporated into the soil either by mechanical mixing of the soil with implements or by "watering in" by rainfall or irrigation. The soil of a plant locus may also be treated with antidote by application of the antidote in a dispersible-concentrate form such as a granule. The granule may be applied to a furrow which is prepared for receipt of the crop seed and the herbicide may be applied to the plant locus either before or after in-furrow placement of the antidote-containing granule so that the herbicide and antidote form a "combination". Crop seed may be treated or coated with the antidote compound either while the crop seed is in-furrow just after seeding or, more commonly, the

crop seed may be treated or coated with antidote prior to seeding into a furrow. The herbicide may be applied to the soil plant locus before or after seeding and a "combination" is made when both herbicide and an antidote-coated seed are in the soil.

5 In a preferred method of the invention (a) the herbicidal benzoylisoxazole and/or dione derivative and (b) antidote are applied separately such that the antidote contacts the seed or plant being treated before the herbicidal compound. Separate application can be achieved, for example, by dressing seed with antidote. The seed may also be treated, for example, at the time of planting, for example by
10 applying antidote to seed in a seed furrow or to growing medium (for example soil) which then covers the seed. The herbicidal compound may then be applied to the surface of the growing medium (for example soil) or to a layer of soil above the layer containing antidote.

Apparatus suitable for planting seed, for example in a furrow and applying antidote
15 to a layer of soil to cover the seed is generally known in the art. Herbicide may then be applied to the soil surface, for example by spraying a fluid formulation or distributing a solid formulation.

Separate application can also be achieved, for example, by applying the herbicidal compound and antidote simultaneously, preferably in a single composition, which
20 composition permits contact of the antidote with the seed or plant before contact with the herbicidal compound.

For example the composition may comprise a delayed release formulation of the herbicidal compound, for example the antidote in the composition is released substantially immediately and before delayed release of the herbicidal compound.

25 In one embodiment the method of the invention is preferably performed by applying the antidote directly to the seed before planting. This is generally effected by coating a quantity of crop seed with the antidote and thereafter planting the coated seed.

In a further preferred embodiment of the method of the invention, the antidotes are
30 selected from fenchlorazole; CMPI; 4-hydroxy-1-methyl-3-(1-1H-tetrazol-5-ylmethanoyl)-1H-quinolin-2-one; (S)-MBU and dimepiperate.

The amount of a particular benzoylisoxazole and/or dione herbicide to be applied to the plant locus or crop-growing area will depend upon the nature of the weeds, the particular herbicide used, the time of application, the climate and the nature of the
35 crop. Application rates of from about 0.004kg/ha⁻¹ to 5kg/ha⁻¹ herbicide are

generally suitable, with a rate of about 0.01kg ha^{-1} to 2kg ha^{-1} being preferred, and with a rate of 0.005kg ha^{-1} to 0.3kg ha^{-1} being more preferred.

The amount of antidote used in the method of the invention varies according to a number of parameters including the particular antidote employed, the crop to be
5 protected, the amount and rate of herbicide applied, and the edaphic and climatic conditions prevailing. Also, the selection of the specific antidotes for use in the method of the invention, the manner in which it is to be applied and the determination of the activity which is non-phytotoxic but antidotally effective, can be readily performed in accordance with common practice in the art.

10 The antidote is applied in combination with the herbicide in a non-phytotoxic antidotally effective amount. By "non-phytotoxic" is meant an amount of the antidote which causes at most minor or no injury to the desired crop species. By "antidotally-effective" is meant an antidote used in an amount which is effective as an antidote with the herbicide to decrease the extent of injury caused by the
15 herbicide to the desired crop species.

The herbicide/safener combination according to the invention may also be employed for controlling harmful plants in crops of genetically engineered plants which are either known or still to be developed. As a rule, the transgenic plants are distinguished by particular, advantageous properties, for example by resistances to
20 certain crop protection agents, resistances to plant diseases or pathogens causing plant diseases such as particular insects or microorganisms such as fungi, bacteria or viruses. Other particular properties relate for example, to the harvested material in terms of quantity, quality, storing properties, composition and specific constituents. Thus, transgenic plants are known which have an increased starch
25 content or an altered starch quality, or those where the harvested material has a different fatty acid composition.

The use of the combinations according to the invention in economically important transgenic crops of useful plants and ornamentals, for example cereals such as wheat, barley, rye, oats, panic grasses, rice, cassava and maize or else crops of
30 sugar beet, cotton, soya, oilseed rape, potatoes, tomatoes, peas and other types of vegetables.

When the combinations according to the invention are applied in transgenic crops, effects on harmful plants to be observed in other crops are frequently accompanied
35 by effects which are specific for application in the transgenic crop in question, for

example an altered or specifically widened weed spectrum which can be controlled, altered application rates which may be used, preferably good compatibility with the herbicides to which the transgenic crop is resistant, and altered growth and yield of the transgenic crop plants.

- 5 The invention therefore also relates to the use of the combination according to the invention for controlling harmful plants in transgenic crop plants.

The following non-limiting examples illustrate the invention wherein Antidote A is ethyl 5,5-diphenylisoxazoline-3-carboxylate and Antidote B is 5,5-

- 10 diphenylisoxazoline-3-carboxylic acid.

Example 1

Maize seeds were sown in non-sterile loam and safener, dissolved in acetone was applied to the soil surface. After 30 minutes a treatment of herbicide, Compound
15 A [5-cyclopropyl-4-(2-methylsulphonyl-4-trifluoromethylbenzoyl) isoxazole] was applied to the treated soil.

A visual assessment of the percentage phytotoxicity (measured as a reduction in green plant matter or plant height) compared to an untreated control was made 14 days after treatment (DAT).

- 20 Maize seeds were sown in non-sterile loam and grown up to a 1.5 -2 leaf stage. Antidote, dissolved in acetone, was applied post-emergence to the soil surface. After 30 minutes a treatment of herbicide, Compound A [5-cyclopropyl-4-(2-methylsulphonyl-4-trifluoromethylbenzoyl) isoxazole] was applied to the treated soil.

- 25 A visual assessment of the percentage phytotoxicity compared with an untreated control was made 14 DAT.

Pre-emergence activity of Compound A on maize in the presence of antidotes.

	<u>g/ha of antidote</u>	<u>% phytotoxicity</u>
Cpd A	63	15
Cpd A + mefenpyr-diethyl	63	8.75
Cpd A + fenchlorazole-ethyl	63	11.3
Cpd A + Antidote A	63	11.3
Cpd A + Antidote B	63	8.75

Post-emergence activity of Compound A (63g/ha) on maize in the presence of antidotes.

	<u>g/ha of antidote</u>	<u>% phytotoxicity</u>
Cpd A	63	27.5
Cpd A + mefenpyr-diethyl	63	25
Cpd A + fenchlorazole-ethyl	63	27.5
Cpd A + Antidote A	63	5
Cpd A + Antidote B	63	5

5

Post-emergence activity of Compound A (125g/ha) on maize in the presence of antidotes.

	<u>g/h of antidote</u>	<u>% phytotoxicity</u>
Cpd A	125	43
Cpd A + Antidote A	31	15
Cpd A + Antidote A	63	10
Cpd A + Antidote A	125	15
Cpd A + Antidote B	31	19
Cpd A + Antidote B	63	18
Cpd A + Antidote B	125	25

According to a further feature of the present invention, there are provided
 10 herbicidal compositions comprising:

(a) a herbicidally effective amount of a benzoylisoxazole and/or dione derivative of formula (I) or an agriculturally acceptable salt or metal complex thereof; and

(b) an antidotally effective amount of an antidote compound or an agriculturally acceptable salt thereof;

5 in association with, and preferably homogeneously dispersed in, one or more compatible herbicidally- acceptable diluents or carriers and/or surface-active agents (i.e. diluents or carriers or surface-active agents of the type generally acceptable in the art as being suitable for use in herbicidal compositions and which are compatible with the herbicides of the present invention). In preferred compositions
10 the antidote contacts the seed or plant being treated before the herbicidal compound. For example, the herbicidal compound may be in a delayed release composition. The term "homogeneously dispersed" is used to include compositions in which the benzoylisoxazole and/or dione of formula (I) and antidote are dissolved in the other components. The term "herbicidal composition" is used in a
15 broad sense to include not only compositions which are ready for use as herbicides but also concentrates which must be diluted before use.

The ratio of herbicide to antidote may vary depending upon the crop to be protected, weed to be inhibited, herbicide used, etc., but normally an herbicide-to-antidote ratio ranging from 1:25 to 60:1 parts by weight may be
20 employed, although much higher rates of antidote may be used, e.g., 1:100 to 1:300 parts by weight of herbicide to-antidote. The preferred weight ratio of herbicide-to-antidote is from 1:10 to 30:1. Another preferred weight range ratio is from 1:1 to 20:1, with an even more preferred weight ratio range from 2:1 to 15:1.

25 Preferably, the compositions contain from 0.05 to 90% by weight of benzoylisoxazole and/or dione of formula (I) and antidote.

The herbicidal composition may contain solid and liquid carriers and surface-active agents (e.g. wetters, dispersants or emulsifiers alone or in combination). Surface-active agents that may be present in the herbicidal compositions of the present
30 invention may be of the ionic or non-ionic types, for example sulphuricinoleates, quaternary ammonium derivatives, products based on condensates of ethylene oxide with nonyl- or octyl-phenols, or carboxylic acid esters of anhydrosorbitols which have been rendered soluble by etherification of the free hydroxy groups by condensation with ethylene oxide, alkali and alkaline earth metal salts of sulphuric
35 acid esters and sulphonic acids such as dinonyl- and dioctyl-sodium sulphonyl-

succinates and alkali and alkaline earth metal salts of high molecular weight sulphonic acid derivatives such as sodium and calcium lignosulphonates. Examples of suitable solid diluents or carriers are aluminium silicate, talc, calcined magnesia, kieselguhr, tricalcium phosphate, powdered cork, absorbent carbon black and clays such as kaolin and bentonite. Examples of suitable liquid diluents include water, acetophenone, cyclohexanone, isophorone, toluene, xylene, and mineral, animal, and vegetable oils (these diluents may be used alone or in combination).

Herbicidal compositions according to the present invention may also contain, if desired, conventional adjuvants such as adhesives, protective colloids, thickeners, penetrating agents, stabilisers, sequestering agents, anti-caking agents, colouring agents and corrosion inhibitors. These adjuvants may also serve as carriers or diluents.

Granular formulations may be prepared by absorbing the compounds of the present invention (dissolved in suitable solvents, which may, if desired, be volatile) onto the solid diluents or carriers in granular form and, if desired, evaporating the solvents, or by granulating compositions in powder form obtained as described above.

Powders, dispersible granules and liquid compositions in the form of concentrates may be diluted with water or other suitable diluents, for example mineral or vegetable oils, particularly in the case of liquid concentrates in which the diluent or carrier is an oil, to give compositions ready for use.

The wettable powders (or powders for spraying) usually contain from 20 to 95% of combination, and they usually contain, in addition to the solid vehicle, from 0 to 5% of a wetting agent, from 3 to 10% of a dispersant agent and if necessary, from 0 to 10% of one or more stabilisers and/or other additives such as penetrating agents, adhesives or anti-caking agents and colourings.

The aqueous suspension concentrates, which are applicable by spraying, are prepared in such a way as to obtain a stable fluid product (by fine grinding) which does not settle out and they usually contain from 10 to 75% of combination, from 0.5 to 15% of surface acting agents, from 0.1 to 10% of thixotropic agents, from 0 to 10% of suitable additives such as antifoams, corrosion inhibitors, stabilisers, and water or an organic liquid in which the active substance is sparingly soluble or insoluble. Some organic solid substances or inorganic salts can be dissolved in order to assist in preventing sedimentation or as antifreeze for the water.

Application of the herbicide, antidote, or mixture thereof, can be carried out by conventional techniques utilizing, for example, hand-carried or tractor-mounted spreaders, power dusters, boom and hand sprayers, spray dusters, and granular applicators. If desired, application of the compositions of the invention to plants
5 can be accomplished by incorporating the compositions in the soil or other media. Herbicidal compositions according to the present invention may also comprise (a) and (b) in association with, and preferably homogeneously dispersed in, one or more other pesticidally active compounds and, if desired one or more compatible pesticidally acceptable diluents and carriers. Examples of other pesticidally active
10 ingredients include fungicides, insecticides, plant growth regulators and, most preferably, herbicides.


The optional partner herbicides which may be combined with the derivatives of formula (I) and antidote are preferably selected from chloroacetamides (e.g. metolachlor, acetochlor, alachlor), sulfonyleureas, thiocarbamates,
15 dithiocarbamates, metribuzin, sulfentrazone, flumetsulam, clorasulam-methyl, oxasulfuron, flumiclorac, bentazon, chlorimuron, linuron, clomazone, dimethenamid, pendimethalin, trifluralin, clethodim and acifluorfen, bifenox, diflufenican, diuron, atrazine and ametryne.

According to a further feature of the present invention there is provided a product
20 comprising:

- (a) a herbicidally effective amount of a benzoylisoxazole and/or dione derivative of formula (I), or an agriculturally acceptable salt or metal complex thereof; and
 - (b) an antidotally effective amount of an antidote compound or an agriculturally acceptable salt thereof;
- 25 as a combined preparation for separate, simultaneous or sequential use in the control of weeds at a crop locus, the antidote preferably contacting the seed or plant before the herbicidal compound.

CLAIMS


1. A method of reducing phytotoxicity to a crop (especially maize) at a locus caused by the application thereto of a herbicidal benzoylisoxazole and/or dione derivative of formula (I):




(I)

wherein:


A is a group (A-1) to (A-7):




(A-1)




(A-2)




(A-3)




(A-4)



(A-5)




(A-6)




(A-7)

or a corresponding formula (A-6a) or (A-7a):



(A-6a)



(A-7a)

in which the position of the carbonyl group and the group Q are reversed and the double bond in the ring is attached to the carbon atom attached to the group Q;

- 5 R is a hydrogen atom or a halogen atom; a straight- or branched chain alkyl, alkenyl or alkynyl group containing from one to six carbon atoms which is optionally substituted by one or more halogen atoms; a cycloalkyl group containing from 3 to 6 carbon atoms optionally substituted by one or more groups R^5 , one or more halogen atoms or a group $-CO_2R^3$; or a group selected from $-CO_2R^3$, $-COR^5$,
 10 cyano, nitro, $-CONR^3R^4$ and $-S(O)_kR^{13}$;

R^1 is a straight- or branched-chain alkyl, alkenyl or alkynyl group containing up to six carbon atoms which is optionally substituted by one or more halogen atoms; or a cycloalkyl group containing from three to six carbon atoms optionally substituted by one or more groups R^5 or one or more halogen atoms;

- 15 R^2 is a halogen atom; a straight- or branched-chain alkyl, alkenyl or alkynyl group containing up to six carbon atoms which is optionally substituted by one or more halogen atoms; a straight- or branched-chain alkyl group containing up to six carbon atoms which is substituted by one or more groups $-OR^5$; or a group selected from nitro, cyano, $-CO_2R^5$, $-S(O)_pR^6$, $-O(CH_2)_mOR^5$, $-COR^5$, $-NR^{11}R^{12}$,
 20 $-N(R^8)SO_2R^7$, $-N(R^8)CO_2R^7$, $-OR^5$, $-OSO_2R^7$, $-SO_2NR^3R^4$, $-CONR^3R^4$, $-CSNR^3R^4$, $-(CR^9R^{10})_y-S(O)_qR^7$ and $-SF_5$;

- or two groups R^2 , on adjacent carbon atoms of the phenyl ring may, together with the carbon atoms to which they are attached, form a 5 to 7 membered saturated or unsaturated heterocyclic ring containing up to three ring heteroatoms selected
 25 from nitrogen, oxygen and sulfur, which ring is optionally substituted by one or more groups selected from halogen, nitro, $-S(O)_pR^{13}$, C_{1-4} alkyl, C_{1-4} alkoxy, C_{1-4} haloalkyl, C_{1-4} haloalkoxy, $=O$ (or a 5- or 6- membered cyclic acetal thereof),

and =NO-R³, it being understood that a sulphur atom, where present in the ring, may be in the form of a group -SO- or -SO₂-;

z is an integer from one to five: when z is greater than one the groups R² may be the same or different;

5 R³, R⁴ and R¹⁰⁹ are each independently a hydrogen atom, or a straight- or branched chain alkyl group containing up to six carbon atoms which is optionally substituted by one or more halogen atoms;

R⁵ and R¹¹⁰ are each independently a straight- or branched-chain alkyl group containing up to six carbon atoms which is optionally substituted by one or more
10 halogen atoms or a straight- or branched-chain alkenyl or alkynyl group containing from two to six (preferably from three to six) carbon atoms which is optionally substituted by one or more halogen atoms;

R⁶ and R⁷, which may be the same or different, are each R⁵; or phenyl optionally substituted by from one to five groups which may be the same or different
15 selected from a halogen atom, a straight- or branched-chain alkyl group containing up to six carbon atoms which is optionally substituted by one or more halogen atoms, nitro, cyano, -CO₂R⁵, -S(O)_pR¹³, -NR¹¹NR¹², -OR⁵ and -CONR³R⁴;

R⁸, R⁹ and R¹⁰ are each a hydrogen atom or R⁶;

R¹¹ and R¹² are each a hydrogen atom or R⁵;

20 R¹³ and R¹¹¹ are each a straight- or branched-chain alkyl group containing up to six carbon atoms which is optionally substituted by one or more halogen atoms;

Q is hydroxy, C1-6 alkoxy, OR¹¹², SR¹¹² or SR¹¹¹;

L is oxygen or NR¹⁰⁹;

R¹⁴, R^{14a}, R^{14b}, R¹⁵, R^{15a}, R^{15b}, R¹⁶, R^{16a}, R^{16b}, R¹⁰⁰, R¹⁰¹, R¹⁰², R¹⁰³,
25 R¹⁰⁴, R¹⁰⁵, R¹⁰⁶, R¹⁰⁷ and R¹⁰⁸ are each the same or different groups selected from hydrogen, R¹¹⁰, -(CH₂)_uCO₂R¹⁰⁹, halogen, cyano, C1-6 alkoxy, -(CH₂)_x-[phenyl optionally substituted by from one to five groups R¹¹³ which may be the same or different], and cycloalkyl containing from three to six carbon atoms optionally substituted by C1-6 alkyl or -S(O)_pR¹¹¹ ;

30 R¹¹² is phenyl optionally substituted by from one to five groups selected from halogen, C1-6 alkyl, C1-6 haloalkyl, C1-6 alkoxy and nitro;

R¹¹³ is a group selected from halogen, R¹¹⁴, nitro, cyano, -CO₂R¹¹⁵,
-S(O)_pR¹¹¹, -OR¹¹¹ and -NR¹¹⁵R¹¹⁶;

R¹¹⁴ is a straight- or branched- chain alkyl group containing one to three carbon atoms optionally substituted by one or more halogen atoms;

5 R¹¹⁵ and R¹¹⁶ which may be the same or different, are each a hydrogen atom or R¹¹⁰;

p, q and u are each independently zero, one or two;


k and m are each one, two or three;

x is zero or one;

10 y is an integer from one to four; when y is greater than one, the groups R⁹ and R¹⁰ may be the same or different;

or an agriculturally acceptable salt or metal complex thereof; which method comprises applying to the locus of the crop, preferably before the herbicidal compound, an antidotally effective amount of an antidote compound, and optional
15 partner herbicide.

2. A method according to claim 1 in which the isoxazole or dione herbicide has the general formula (Ia):



(Ia)

are those wherein:

R is hydrogen or -CO₂Et;


R¹¹⁷ is selected from -S(O)_pMe, Me, Et, a chlorine, bromine or fluorine atom, methoxy, ethoxy and -CH₂S(O)_qMe;

25 R¹¹⁸ is selected from a hydrogen atom, a chlorine, bromine or fluorine atom, methoxy, ethoxy and -S(O)_pMe;

R¹¹⁹ is selected from a hydrogen atom, a chlorine, bromine or fluorine atom, methoxy and CF₃; and

p and q each independently have the values zero, one or two.


3. A method according to claim 1 or 2 in which the isoxazole or dione herbicide has the general formula (Ib):



(Ib)


wherein R^{120} is chlorine, bromine or trifluoromethyl; and R is hydrogen or $-CO_2Et$.

4. A method according to any one of the preceding claims in which a substituted phenyl ring as defined in formulae (I); (Ia); or (Ib) as depicted in claim 1, 2 or 3 is attached to a grouping;



5. A method according to claim 4 in which the phenyl ring is substituted by two groups independently selected from halogen, alkyl, $S(O)_p$ alkyl ($p = 0, 1$ or 2) or haloalkyl.

6. A method according to claim 1, 2 or 3 in which a substituted phenyl ring as defined above in formula (I); (Ia); or (Ib) is attached to a grouping;



7. A method according to claim 1 wherein the compound of formula (I) is:
 5-cyclopropyl-4-[2-chloro-3-ethoxy-4-(ethylsulphonyl)benzoyl]isoxazole;
 4-(4-chloro-2-methylsulphonylbenzoyl)-5-cyclopropylisoxazole;
 5-cyclopropyl-4-(2-methylsulphonyl-4-trifluoromethylbenzoyl)isoxazole;
 4-(4-bromo-2-methylsulphonylbenzoyl)-5-cyclopropylisoxazole;
 5-cyclopropyl-4-[4-fluoro-3-methoxy-2-(methylsulphonyl)benzoyl]isoxazole;

4-(4-bromo-2-methylsulphonylmethylbenzoyl)-5-cyclopropylisoxazole;
ethyl 5-cyclopropyl-4-(2-methylsulphonyl-4-trifluoromethylbenzoyl) isoxazole-3-
carboxylate;
2-[2-chloro-(4-methylsulphonyl)benzoyl]-1,3-cyclohexanedione;
5 2-[2-nitro-(4-methylsulphonyl)benzoyl]-1,3-cyclohexanedione;
2-(2,3-dihydro-5,8-dimethyl-1,1-dioxospiro[4H-1-benzothiine-4,2' [1,3]dioxolan]-6-
ylcarbonyl)cyclohexane-1,3-dione;
5-cyclopropyl-4-(2-methylsulphonyl-4-trifluoromethylbenzoyl)-3-methylthio-
isoxazole; and
10 2-cyano-3-cyclopropyl-1-(2-methylsulphonyl-4-trifluoromethylphenyl) propan-1,3-
dione.

8. A method according to claim 7 in which the compound is
5-cyclopropyl-4-(2-methylsulphonyl-4-trifluoromethylbenzoyl)isoxazole or 2-[2-nitro-
(4-methylsulphonyl)benzoyl]-1,3-cyclohexanedione.

15 9. A method according to claim 1 in which the antidote is selected from:
flurazole; fenchlorazole-ethyl; fenchlorazole; benoxacor; dichlormid; fenclorim;
furilazole; mefenpyr-diethyl; CMPI; 4-hydroxy-1-methyl-3-(1-1*H*-tetrazol-5-
ylmethanoyl)-1*H*-quinolin-2-one; daimuron; (S)-MBU; dimepiperate; 5,5-
diphenylisoxazoline-3-carboxylic acid; and ethyl 5,5-diphenylisoxazoline-3-
20 carboxylate.

10. A method according to claim 1 in which the antidote is selected from
fenchlorazole; CMPI; 4-hydroxy-1-methyl-3-(1-1*H*-tetrazol-5-ylmethanoyl)-1*H*-
quinolin-2-one; (S)-MBU and dimepiperate.

11. A method according to any one of the preceding claims in which the crop
25 plant to be protected is maize.

12. A method according to claim 1 in which the application rate of the
benzoylisoxazole and/or dione of formula (I) is from 0.004kg to 5kg per hectare.

13. A method according to claim 1 in which the application rate of the
benzoylisoxazole and/or dione of formula (I) is from 0.01kg to 2kg per hectare.

30 14. A method according to any one of the preceding claims in which (a) the
herbicidal benzoylisoxazole and/or dione derivative and (b) antidote are applied
separately such that the antidote contacts the seed or plant being treated before
the herbicidal compound.

15. A herbicidal composition comprising:

- (a) a herbicidally effective amount of a benzoylisoxazole and/or dione derivative of formula (I) or an agriculturally acceptable salt or metal complex thereof, optionally in combination with a partner herbicide; and
- (b) an antidotally effective amount of an antidote compound;
- 5 in association with a herbicidally acceptable diluent or carrier and/or surface active agent.
16. A composition according to claim 15 which comprises the component (a) as a delayed release formulation.
17. A composition according to claim 15 or 16 in which the weight ratio of the
10 compound of formula (I):antidote is from 1:25 to 60:1.
18. A product comprising:
- (a) a herbicidally effective amount of a benzoylisoxazole and/or dione derivative of formula (I), or an agriculturally acceptable salt or metal complex thereof; and
- (b) an antidotally effective amount of an antidote;
- 15 wherein said antidote is antidotally effective to said benzoylisoxazole and/or dione derivative;
- as a combined preparation for separate, simultaneous or sequential use in the control of weeds at a locus.
19. A product according to claim 18 as a combined preparation for use in which
20 the antidote contacts the seed or plant being treated before the herbicidal compound.
20. A method according to claim 1 substantially as hereinbefore described.
21. A composition according to claim 15 substantially as hereinbefore described.
- 25 22. A product according to claim 18 substantially as hereinbefore described.

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
15 March 2001 (15.03.2001)

PCT

(10) International Publication Number
WO 01/17350 A1

(51) International Patent Classification⁷: A01N 43/80,
41/10, 43/90 // (A01N 43/80, 43:80, 43:653, 43:56, 25:32)
(A01N 41/10, 43:80, 43:653, 43:56, 25:32) (A01N 43/90,
43:80, 43:653, 43:56, 25:32)

(21) International Application Number: PCT/EP00/09339

(22) International Filing Date:
8 September 2000 (08.09.2000)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
9921220-1 8 September 1999 (08.09.1999) GB
0012090-7 19 May 2000 (19.05.2000) GB

(71) Applicant (for all designated States except US): AVENTIS
CROPSCIENCE UK LIMITED [GB/GB]; Fyfield Road,
Ongar, Essex CM5 0HW (GB).

(72) Inventors; and

(75) Inventors/Applicants (for US only): PALLETT, Ken
[GB/GB]; Aventis CropScience UK Ltd, Fyfield Road, On-
gar, Essex CM5 0HW (GB). SLATER, Ashley [GB/GB];
Aventis CropScience UK Ltd, Fyfield Road, Ongar, Essex
CM5 0HW (GB).

(74) Agent: AVENTIS CROPSCIENCE SA; Département
Propriété Industrielle - N° 153, B.P. 9163, F-69263 Lyon
Cedex 09 (FR).

(81) Designated States (national): AE, AG, AL, AM, AT, AU,
AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CR, CU, CZ,
DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR,
HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR,
LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ,
NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM,
TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW.

(84) Designated States (regional): ARIPO patent (GH, GM,
KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian
patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European
patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE,
IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG,
CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).

Published:

- With international search report
- Before the expiration of the time limit for amending the
claims and to be republished in the event of receipt of
amendments.

For two-letter codes and other abbreviations, refer to the "Guid-
ance Notes on Codes and Abbreviations" appearing at the begin-
ning of each regular issue of the PCT Gazette.

(54) Title: NEW HERBICIDAL COMPOSITIONS

(57) Abstract: The present invention provides a method of reducing phytotoxicity to crops (especially maize) caused by a herbicidal benzoylisoxazole and/or dione derivative of formula (I) or an agriculturally acceptable salt or metal complex thereof; which method comprises applying to the locus of the crop an antidotally effective amount of an antidote compound, optionally with a partner herbicide.



WO 01/17350 A1

COMBINED DECLARATION FOR PATENT APPLICATION AND POWER OF ATTORNEY

As below named inventor, I hereby declare that:

My residence, post office address and citizenship are as stated below, I believe I am the original, first and sole inventor (if only one name is listed below) or an original, first and joint inventor (if plural names are listed below) of the subject matter which is claimed and for which a patent is sought on the invention entitled

New Herbicidal Compositions

the specification of which

- is attached hereto

- was filed on September 8, 2000 as International Application PCT/EP00/09339 and including all the amendments through the date hereof.

I hereby state that I have reviewed and understand the contents of the above identified specification, including the claims, as amended by any amendment referred to above.

I acknowledge the duty to disclose to the Office all information known to me to be material to patentability as defined in Title 37, Code of Federal Regulations, §1.56.

I hereby claim foreign priority benefits under Title 35, United States Code, §119 of any foreign application(s) for patent or inventor's certificate listed below and have also identified below any foreign application for patent or inventor's certificate having a filing date before that of the application on which priority is claimed:

Prior Foreign Application(s) for which Priority is Claimed:

GB 9921220-1 of September 8, 1999

GB 0012090-7 of May 19, 2000

And I hereby appoint

William F. Lawrence, Registration No. 28,029, of the firm FROMMER LAWRENCE & HAUG, LLP whose post office address is 745 Fifth Avenue, New York, New York 10151, or their duly appointed associate, my attorneys, with full power of substitution and revocation, to prosecute this application, to make alterations and amendments therein, to file continuation and divisional applications thereof, to receive the Patent, and to transact all business in the Patent and Trademark Office and in the Courts in connection therewith, and specify that all communications about the application are to be directed to the following correspondence address:

William F. Lawrence, Esq.
c/o FROMMER, LAWRENCE & HAUG LLP
745 Fifth Avenue
New York, New York 10151

Direct all telephone calls
to: (212) 588-0800, to the
attention of:
William F. Lawrence

I hereby declare that all statements made herein of my own knowledge are true and that all statements made on information and belief are believed to be true; and further that these statements were made with the knowledge that willful false statements and the like so made are punishable by fine or imprisonment, or both, under Section 1001 of Title 18 of the United States Code and that such willful false statements may jeopardize the validity of the application or any patent issued thereon.

INVENTOR(S) / Residence

1) Ken PALLETT, Adelheidstraße 10, 61462 Königstein, Germany *DEX*Signature: Date: 14/01/022) Ashley SLATER, Aventis CropScience UK Ltd., Fyfield Road, Ongar, Essex CM5 0HW, Great BritainSignature: Date: 14/01/02

The inventors are citizens of Great Britain.

Post Office Address of all Inventors:

Aventis CropScience GmbH
Patent- und Lizenzabteilung
Industriepark Höchst
Bldg. K 801
D-65926 Frankfurt
Germany